Microbial Menaces and their Biodiversity Present in Crops Irrigated with Polluted Yamuna Water in Peri-urban Agriculture area of Delhi and National Capital regions, India

Sumit Pal^{1,2}, Neelam Patel¹, Anushree Malik² and D.K. Singh³

¹Water Technology Centre, Indian Agricultural Research Institute, New Delhi - 110 012, India. ²CRDT, Indian Institute of Technology, New Delhi, India. ³Department of Zoology, University of Delhi, New Delhi, India.

https://doi.org/10.22207/JPAM.10.3.50

(Received: 24 May 2016; accepted: 03 July 2016)

Crops can become contaminated with microorganisms capable of causing human diseases while still on the plant in fields or orchards, or during harvesting. The objective of this study was to assess the contamination level of microbial load, yeasts, molds, Coliform and Escherichia coli on their surface and biodiversity indexes of microorganisms. The bacterial counts were found to be higher than the Yeast and Moulds counts. The range of bacterial population is 8.72-5.04 log cfu/g and for the fungal is 6.53 - 2.30 log cfu/g. Organisms isolated from vegetables samples included Saccharomyces, Aspergillus, Rhizopus, Staphylococcus, Klebsiella, Streptococcus, Bacillus, Escherichia, Pseudomonas, Salmonella and Diplococcus with various percentages of occurrences. Streptococcus and Aspergillus are the most frequently isolated genera in all studied crops. Staphylococcus (86.36%) and Pseudomonas (77.27%) present in the studied crops. It's the main cause of unwanted diseases in Delhi and NCR. The biodiversity indexes i.e. Simpson (0.20) and Shannon (3.31) of microorganism show the high species richness and diversified microbial community. The microorganisms isolated are opportunistic and mostly pathogenic in nature. The microbial load also very high which gives the signal of health alertness before using these contaminated vegetables.

Keywords: Aspergillus, Diplococcus, Microbial load, Streptococcus, Vegetables, Simpson Index.

The consumption of contaminated raw vegetables may lead to food poisoning due to presence of harmful and pathogenic microorganism¹ The outbreaks of food poisoning cases have therefore highlighted the importance of microbial quality assessment of food commodities. Raw vegetables are known potential for a wide range of microorganisms, including human pathogens². Pathogens commonly detected in fresh vegetables are *E. coli*, coliform bacteria, *Staphylococcus* sp., *Salmonella* sp *Campylobacter spp.*, *Shigella spp.*, *Clostridium*

botulinum, Listeria monocytogenes, Clostridium perfringens, some Bacillus spp., and Streptococcus³ The increasing numbers of pathogens in vegetables are due to the widespread use of human and animal faecal waste water and Industrial waste water in agricultural practices⁴. It is not surprising that enteric pathogens can contaminate agricultural produce and cause outbreaks of illness due to the following contaminated consumption⁵. Spores of Clostridium botulinum, Clostridium perfringens, and Bacillus cereus can also be isolated from fresh vegetables and Listeria monocytogenes can be found in decaying vegetables⁶. These microorganisms are likely to contaminate raw fruits and vegetables through contact with faeces, sewage, untreated irrigation water or surface water⁷ Factors contribute

E-mail: sumit microbe@gmail.com

^{*} To whom all correspondence should be addressed. Tel.: 9654647834:

to real increases in diseases associated with fruits and vegetables due to use of wastewater, increased application of improperly composted manures to soils in which fruits and vegetables are grown, extended time between harvesting and consumption, and changing food consumption patterns8 The presence of a pathogen on produce is of less consequence if the rind, skin or peel is to be removed before consumption. Potato, Onion, Bottle gourd, Bitter gourd and all other vegetables having the peel, fall in this category9 However, Microorganisms that have become trapped on the inner leaves of certain vegetables can be particularly difficult to remove by routine cleansing practices¹⁰. Application of improperly composted manure or water containing raw sewage to fields, or the use of water contaminated with faeces during processing of fruits and vegetables must be avoided¹¹. The microbial quality of vegetables grown with wastewater is highly alarming. Several studies throughout the world have demonstrated a very close relation between the consumption of fruits and vegetables irrigated with raw wastewater and many food borne diseases like gastroenteritis, cholera, chemical toxicity etc12

Due to increasing consumption of vegetables as a result of awareness of its health benefits and report showing impacts of vegetables irrigated by using untreated wastewater on health were available¹³. The need to assess the microbial quality of irrigated vegetables in Delhi, India is eminent in order to propose alternatives to prevent consumers from disease outbreaks. Therefore, the present study was undertaken to examine the quality of Crops for microbial contamination, irrigated from waste water from Delhi and peri urban

MATERIALS AND METHOD

Study area

The study was carried out in Delhi –NCR peri urban area. The samples of crops were collected from the 15 different sites of Yamuna pusta region. i.e. Palla, Christian ashram, Jagatpur, Sonia Vihar, Wazirabad, Shastri park, Indraprastha, Okhla, Noida, Basantpur, Nehru vihar, Daryia nalla, Punjabi bagh, Keshopura and Nilothi. Crops grown in these areas are gone for the sale in local market. [Fig. 1].



Fig. 1. Map of 15 sampling sites of Delhi-NCR peri urban area

J PURE APPL MICROBIO, 10(3), SEPTEMBER 2016.

Sample collection and processing

Four samples of each crop from 15 Different sites were collected as shown in Fig. 1. Random sampling procedure was adopted to collect the sample materials. The vegetables samples were collected in sterile polythene zip bags to laboratory for storage in refrigerator.

Microbial analysis of crops

Twenty five gram [25 g] of each crop sample were weighed and blended in 100 mL of sterile saline solution for 2 minutes under sterile conditions. The blender was carefully disinfected to prevent any cross contamination. The homogenates were collected in sterile bottles and stored at -20°C until needed. Aliquots [1 mL] of each homogenate were serially diluted in sterile saline solution. The diluents of buffered peptone water were then inoculated on to the respective media. Yeast and Mould count in crops was determined by the procedure described by IS 5403: 1999¹⁴ Coliforms bacteria in crops were determined by the procedure described by IS 5401[II]: 2002¹⁵ Total plate count in crops was determined by the procedure described by IS 5402:200216 E. coli in crops was determined by the procedure described by IS 5887[I]: 1976, 17 Salmonella, Streptococcus and Staphylococcus by IS 5887: 1976, ¹⁸Vibrio by IS5887:1976,18Pseudomonas IS 13428:2005.19 Gram reaction and biochemical tests such as catalyse, oxidase, urease, citrate, methyl red test, Voges Proskauer, indole, sugar fermentation were performed to identify other bacteria [20] while fungal isolates were determined

based on their cultural and microscopic morphology²¹

RESULTS AND DISCUSSION

Total twenty two different varieties of crops studied for the Microbial quality analysis. After studying all the crop samples, the bacterial count as found more than the fungal count in studied crops [Table 1]. The microbial study of crops also shows that the different crops have different concentration of microbial load on it which makes the crops, specific for the contamination and also not suitable to eat in regards to human.

The organisms isolated from the 22 different varieties of samples are Saccharomyces, Aspergillus, Rhizopus, Staphylococcus, Klebsiella, Streptococcus, Bacillus, Escherichia, Pseudomonas, Salmonella and Diplococcus. The percentage of occurrence of Aspergillus and Streptococcus (100), Staphylococcus(86.6), P s e u d o m o n a s (77.27), E.coli(59.09), Salmonella(36.36), Bacillus(36.6), Saccharomyces(31.81), Rhizopus(27.27), Klebsiella(22.72) and Diplococcus(4.54) in studied crops. The bacterial and fungal count of microorganism ranges from 8.72-5.08 log cfu/g and 6.53-2.3 log cfu/g respectively [Table 1].

On studying, the percentage of occurrence of microorganism in the crops, Streptococcus and Aspergillus are the organism present in all crops. While diplococcus is rarest




Fig. 2. Microbial population in crops

microorganism in the studied crops. Most of the microorganisms are pathogenic, which cause serious diseases to human who feed on these contaminated crops. *Pseudomonas* infects the

pulmonary tract, urinary tract, burns, wounds, and also causes other blood infections²². *Aspergillus* causes otomycosis and aspergillosis in human²³. *Bacillus* causes anthrax and food poisoning²⁴.




Fig. 3. Biodiversity Indexes

Table 1. Microbial load [fungus and bacteria] on crops

S.No.	Crops	Fungal Count [log cfu/g]±SD	Bacterial Count [log cfu/g] ±SD
1	Cucumber	4 ± 0.44	6.4 ± 0.45
2	Rice	2.3 ± 0.09	5.07 ± 1
3	Bitter gourd	3.3 ± 0.18	7.6 ± 0.31
4	Lady finger	$4.4 \hspace{0.1cm} \pm \hspace{0.1cm} 0.21$	7.8 ± 0.39
5	Radish	$4.1 \hspace{0.1cm} \pm \hspace{0.1cm} 0.06$	7.5 ± 0.12
6	Ridge gourd	5.3 ± 0.08	7.6 ± 0.39
7	Red Spinach	$4.6 \hspace{0.1cm} \pm \hspace{0.1cm} 0.04$	7.8 ± 0.35
8	Mustard	$4.4 \hspace{0.1cm} \pm \hspace{0.1cm} 0.11$	7.8 ± 0.24
9	Wheat	2.7 ± 0.15	5.04 ± 0.07
10	Spinach	5.7 ± 0.99	7.9 ± 0.10
11	Carrot	$4.1 \hspace{0.1cm} \pm \hspace{0.1cm} 0.22$	7.5 ± 0.30
12	Armenian Cucumber	4 ± 0.04	7.5 ± 0.23
13	Pumpkin	3.3 ± 0.32	7.3 ± 0.15
14	Bottle gourd	5.4 ± 0.17	7.7 ± 0.17
15	Sorgham	4 ± 0.03	7.3 ± 0.06
16	Beans	3.3 ± 0.03	7.3 ± 0.06
17	Eddoe	6.5 ± 0.29	7.7 ± 0.06
18	Cabbage	$4.6 \hspace{0.1cm} \pm \hspace{0.1cm} 0.25$	7.3 ± 0.22
19	Pigweed	$4.08~\pm~0.08$	$7.5 ~\pm~ 0.40$
20	Brinjal	3.3 ± 0.19	7.2 ± 0.02
21	Cauliflower	$4.5 \pm \ 0.07$	8.7 ± 0.19
22	Tomato	$4.3 \pm \ 0.12$	6.6 ± 0.33

Recommended limits in crops (WHO-ICMSF): log 3 CFU/ g

Diplococcus causes Pneumonia²⁵ Escherichia causes urinary tract infection with diarrhoea and dysentery condition²⁶. Klebsiella causes pneumonia,²⁷urinary infections, septicemia, meningitis, diarrhea and soft tissue infections²⁸. Staphylococcus causes a wide variety of diseases in humans and animals through either toxin production or penetration. Staphylococcal toxins are a common cause of food poisoning, as they can be produced by bacteria growing in improperly stored food items²⁹. The most common sialadenitis is caused by staphylococci, as bacterial infections. Streptococcus causes the diseases that may be caused include streptococcal toxic shock syndrome, necrotizing fasciitis, pneumonia, bacteremia, acute rheumatic fever and acute glomerulonephritis ³⁰. Salmonella causes typhoid fever, paratyphoid fever, and food poisoning.

On other hand, the biodiversity study i.e. Alpha, Beta and Gamma diversity studied in the population [Tables 2 and 3]. The Simpson index comes around 0.20 and the approximation Simpson comes 0.10, the study reveals that the value of Simpson lean towards zero denotes the more diversified community³¹. Shannon index comes 3.31

Table 2. Alpha Biodiversity Indexes

Alpha Diversity	
Simpson Index	0.20
Simpson Index Approximation	0.10
Dominance Index	0.79
Dominance Index Approximation	0.89
Reciprocal Simpson Index	4.97
Alternate Reciprocal Simpson Index	9.94
Shannon Index	3.31
Berger-Parker Dominance Index	0.10
Shannon Index	2.29
Inverted Berger-Parker Dominance Index	9.53
Shannon Index	-0.998
Margalef Richness Index	0.424
Menhinick Index	8.33E-5
Rényi Entropy/Hill Numbers [r=0,1,2,"]	11,9.971,
	9.94,H"-"
Buzas and Gibson's Index	0.906
Gini Coeffificient	6.098
Equitability Index	0.959
ln[] of Hill Numbers [0,1,2,"]:	2.39,2.299,
	2.297,H"·"

which denote that the studied samples are highly diversified. All other values studied in alpha, beta and gamma diversity, support the diversification [Fig. 3].

The microbial quality of crops was investigated. Rhizopus, Aspergillus, Saccharomyces were fungi isolated from crops. Bacteria isolated included Bacillus, Escherichia, Klebsiella, Staphylococcus, Streptococcus, Salmonella, Diplococcus and Pseudomonas with various percentages of occurrences. Earlier, Similar organisms were isolated from fruits or crops³². Fungi are known to cause spoilage of crops. These microorganisms could contaminate the crops through the use of manure in the farm, spores from the atmosphere, wastewater use in irrigation and washing the crops³³. Food-borne pathogens commonly detected in fresh crops which include Salmonella, Shigella, E. coli, Streptococcus and coliform bacteria³⁴. The bacterial and fungal count of microorganism ranges from 8.72-5.08 log cfu/g and 6.53-2.3 log cfu/g, respectively. The presence of these microorganisms in crops creates a public health concern. The Alpha, Beta and Gamma diversity of the studied microorganism shows the excellent Simpson and Shannon index values. The value of these indexes is towards zero which proves that the biodiversity level of microorganism is diversified in studied crops. The high species diversification suggest that a greater number of successful species35 and a more stable ecosystem, more ecological niches are available and the

 Table 3. Beta Biodiversity Indexes

Beta Biodiversity	
Absolute beta Value $[[S_0-c]-[S_1-c]]$:	10
Whittaker's Index [S/alpha]:	1
Sørensen's similarity index:	1
Alternate Whittaker's Index [S/alpha-1]:	0
Sørensen's similarity index [%]:	100%
Jaccard Index:	-1
Routledge beta-R Index:	3.667
Jaccard Index [%]:	-100%
Mountford Index:	-0.22
Number of Common Species:	11
Mountford Index [%]:	-22.22%
Bray Curtis dissimilarity	0
Gamma Biodiversity	
Absolute gamma $[S_0 + S_1 - c]$:	0

environment is less likely to be hostile, complex food webs and environmental change is less likely to be damaging to the ecosystem as a whole.

In the present paper, the authors assessed the microbial load of twenty two different vegetable samples from fifteen sites of Delhi peri urban area. Contamination was found in all the studied crops. Cauliflowers, Lady Finger, Mustard, Red Spinach were the highly contaminated crops. Along with this, the value of Simpson and Shannon index is towards zero which proves that the biodiversity level of microorganism is diversified in studied crops. Beside soil microbes, other sources of their contamination are improper handling and waste water used for irrigation. Wastewater uses in agriculture will increase the microbial load in the crops³⁶. Vegetables may be contaminated with pathogenic microorganisms during growing in the field or during harvesting, post harvesting, handling, processing and distribution. Therefore, vegetables may act as a reservoir for many microorganisms from which they will be colonized inside these vegetables and infect susceptible host. Sanitary measures should be adopted while handling crops to limit the level of microbial contamination

ACKNOWLEDGEMENT

This research was supported by the grants from the National fund for Basic, Strategic and frontier application research in Agriculture, New Delhi, Indian Council of Agricultural Research, New Delhi, India. We acknowledge the excellent assistance of Ram Shankar, Saif Khan, Gaurav Singh, Giriraj Sharma in helping for the field and laboratory assistance.

REFERENCES

- 1. Sivapalasingam S, Friedman CR, Cohen L, Tauxe RV Fresh produce: a growing cause of outbreaks of foodborne illness in the United States, 1973 through 1997. *Journal of Food Protection*®, 2004, **67**(10):2342-2353.
- Leff, J. W., & Fierer, N., Bacterial communities associated with the surfaces of fresh fruits and vegetables. *PloS one*, 2013, 8(3), e59310.
- 3. Jay JM, Loessner M.J, Golden DA *Modern* food microbiology. Springer. 2005.
- 4. Conway, G. R., & Pretty, J. N., Unwelcome

- harvest: agriculture and pollution. Routledge.
 Tauxe RV: Emerging foodborne
- pathogens. International journal of food microbiology, 2013,78(1): 31-41. (2002)
- 6. Siroli, L., Patrignani, F., Serrazanetti, D. I., Gardini, F., & Lanciotti, R., Innovative strategies based on the use of bio-control agents to improve the safety, shelf-life and quality of minimally processed fruits and vegetables. *Trends in Food Science & Technology*, 2014, 46(2), 302-310.
- 7. Said, D. E. S., Detection of parasites in commonly consumed raw vegetables. *Alexandria Journal of Medicine*, 2012, **48**(4), 345-352.
- León JS, Jaykus LA, Moe CL Food safety issues and the microbiology of fruits and vegetables. Microbiologically Safe Foods, 2009, 255-281
- 9. Swamy KRM, Sadashiva AT Tropical Vegetable Cropss. *Biodiversity in Horticultural Cropss*, 2009, 109.
- Ijabadeniyi, O. A., & Buys, E. M., Irrigation water and microbiological safety of fresh produce: South Africa as a case study: a review 2012.
- 11. Selma MV, Allende A, López Gálvez F, Elizaquivel P, Aznar R, Gil MI Potential microbial risk factors related to soil amendments and irrigation water of potato cropss. *Journal of applied microbiology*, 2007, **103**(6): 2542-2549.
- 12. Chalkiadaki O, Dassenakis M, Simantiris NL. Cadmium and Lead Accumulation and Metallothionein overexpression in Tissues of Venus verrucosa Exposed to Contaminated Seawater. In17th International Symposium on Environmental Pollution and its Impact on Life in the Mediterranean Region, 2016, (p. 84).
- Lu Y, Song S, Wang R, Liu Z, Meng J, Sweetman AJ, Jenkins A, Ferrier RC, Li H, Luo W, Wang T. Impacts of soil and water pollution on food safety and health risks in China. Environment international., 2015;77:5-15.
- 14. Chen MM, Liu QH, Xin GR, Zhang JG. Characteristics of lactic acid bacteria isolates and their inoculating effects on the silage fermentation at high temperature. Letters in applied microbiology., 2013; 56(1):71-8.
- IS 5403 Method for Yeast and Mould Count of Foodstuffs and Animal Feeds. 1999.
- 16. IS 5401 (Pt-1) Microbiology-General Guidance for the enumeration of Coliforms. 2002.
- 17. IS 5402 Microbiology- General Guidance for the enumeration of Microorganisms-Colony Cont Technique at 30°C. 2002.
- 18. IS 5887 (Pt-1) Isolation, Identification and

- Enumeration of Escherichia coli. 1976.
- IS 13428 Detection and Enumeration of Pseudomonas aeruginosa (Annexure-D) 2005.
- de Boer E, Beumer RR Methodology for detection and typing of foodborne microorganisms. *International journal of food* microbiology, 1999, 50(1): 119-130.
- Lorch, J. M., Lindner, D. L., Gargas, A., Muller, L. K., Minnis, A. M., & Blehert, D. S., A culturebased survey of fungi in soil from bat hibernacula in the eastern United States and its implications for detection of Geomyces destructans, the causal agent of bat white-nose syndrome. Mycologia, 2012, 12-207.
- Sahu, M. C., Dubey, D., Rath, S., Debata, N. K., & Padhy, R. N., Multidrug resistance of Pseudomonas aeruginosa as known from surveillance of nosocomial and community infections in an Indian teaching hospital. *Journal of Public Health*, 2012, 20(4), 413-423.
- Szigeti, G., Kocsubé, S., Dóczi, I., Bereczki, L., Vágvölgyi, C., & Varga, J., Molecular identification and antifungal susceptibilities of black Aspergillus isolates from otomycosis cases in Hungary. *Mycopathologia*, 2012, 174(2), 143-147
- 24. Granum, P. E., Bacillus cereus Food. *Food Associated Pathogens*, 2013, **20**.
- 25. Morens DM, Taubenberger JK, Fauci AS Predominant role of bacterial pneumonia as a cause of death in pandemic influenza: implications for pandemic influenza preparedness. *Journal of Infectious Diseases*, 2002, **198**(7):962-970.
- Mainil, J., Escherichia coli virulence factors. Veterinary immunology and immunopathology, 2013, 152(1), 2-12.
- 27. Siu, L. K., Yeh, K. M., Lin, J. C., Fung, C. P., & Chang, F. Y., Klebsiella pneumoniae liver abscess: a new invasive syndrome. *The Lancet infectious diseases*, 2012, **12**(11), 881-887.

- Bereket, W., Hemalatha, K., Getenet, B., Wondwossen, T., Solomon, A., Zeynudin, A., & Kannan, S., Update on bacterial nosocomial infections. Eur Rev Med Pharmacol Sci, 2012, 16(8), 1039-1044.
- 29. Le Loir Y, Baron F, Gautier M.: Staphylococcus aureus and food poisoning. *Genet Mol Res*, 2003, **2**(1):63-76.
- 30. Ralph, A. P., & Carapetis, J. R., Group A streptococcal diseases and their global burden. In *Host-Pathogen Interactions in Streptococcal Diseases*, 2012, (pp. 1-27). Springer Berlin Heidelberg.
- 31. Han, D., Ha, H. K., Hwang, C. Y., Lee, B. Y., Hur, H. G., & Lee, Y. K., Bacterial communities along stratified water columns at the Chukchi Borderland in the western Arctic Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 2015, **120**, 52-60.
- 32. Nwachukwu E, Ezeama C, Ezeanya B Microbiology of polyethylene packaged sliced watermelon sold by street vendors in Nigeria. *Afr. J. Microbiol. Res.*, 2008, **2**:192 195.
- Ray B, Bhunia A Fundamental food microbiology 4th Edition, CRC Press USA,2007, p. 492.
- Tambekar D, Mundhada H Bacteriological quality of salad vegetable sold in Amravati city.
 J. Biol. Sci. ,2006, 6(1):28 30.
- Magurran, A. E., Ecological diversity and its measurement. Springer Science & Business Media., 2013.
- Tripathi VK ,Rajput T, Patel N, Quality of municipal wastewater for micro irrigation. *Micro Irriga for Trees an*, 2014, 95.

© The Author(s) 2016. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.