

## Characterization of Methanolic Extract of Red Pigment from *Penicillium purpurogenum* and its Antioxidant Activity

C. Padmapriya\* and R. Murugesan

Department of Agricultural Microbiology  
Tamil Nadu Agricultural University, Coimbatore - 641003, Tamil Nadu, India.

(Received: 30 November 2015; accepted: 10 January 2016)

The fungal pigments are a good alternative to currently used synthetic colourants and / or natural colourants derived from plant materials. An extracellular red pigment producing fungus, *Penicillium purpurogenum* was isolated from soil collected from Parambikulam Tiger Reserve Forest, Kerala. In the present study antioxidant assays viz., Lipid peroxidation, hydroxyl radical and Nitric oxide radical scavenging assays were carried out with fungal red pigment. The pigment extracted from this fungus showed maximum inhibition of lipid peroxidation (30.15 %), hydroxyl radical (74.92 %) and nitric oxide radical scavenging activity (43.23%) at 20 mg ml<sup>-1</sup> of pigment concentration. The red pigment from *P. purpurogenum* was separated by (TLC) yielding three major fractions, viz., Pinkish red, orange and yellow fractions. These fractions were further identified through Gas chromatography and Mass Spectrometry (GC-MS). The identification of the structure of the red pigment was detected using FT-IR spectra, and indicated that the presence of phenolic and quinone compounds and has broad stretching OH, C=C and C-H groups of the aromatic ring.

**Keywords:** Red pigment, *Penicillium purpurogenum*, antioxidant activity, FT-IR, natural food colourants.

The use of natural dyes in food has increased recently due to the marketing advantages with the development of natural ingredients and the consumer concern about the harmful effects of synthetic pigments on health<sup>1</sup>. Pigments are derived from natural sources such as plants, insects, and microorganism. There has been much interest in the development of new natural colourants for use in the food industry owing to strong consumer demand for more natural products. The production of many existing natural colourants of plant origin has a disadvantage of dependence on the supply of raw materials, which are influenced by agro-climatic conditions – in addition, their chemical profile may vary from

batch-to-batch. Moreover, many of the pigments derived from the contemporary sources are sensitive to heat, light, and oxygen, and some may even change their colour in response to pH changes as in case of anthocyanins<sup>2</sup>. Many ascomycetous fungi naturally synthesize and secrete pigments and may thus provide a more reliable source for natural, “organic” food colourants with improved functionalities<sup>3</sup>. The diversity of fungal pigments is not only found in their chemical structures but also in the colour range of these pigments that may add new or additional hues to the colour palette of the existing colourants derived from contemporary sources<sup>4</sup>.

Reactive oxygen species (ROS) are formed as a result of normal metabolic activity (oxidative stress) and from exogenous sources. ROS formed often results in causing cellular and subcellular damage by preoxidation of membrane lipids, denaturation of DNA strands and cellular

\* To whom all correspondence should be addressed.  
E-mail: agri.padma@gmail.com

proteins. In multicellular organisms, cell protects itself from the damage caused by the ROS system by various enzymatic pathways which includes catalase, glutathione peroxidase, glutathione reductase, superoxide dismutase, lipid peroxidase and non enzymatically by beta carotene, vitamin A, C, E<sup>5</sup>. To avoid oxidative stress, antioxidants can play an important role conferring beneficial healthy effects<sup>6</sup>. High dietary intake of proven antioxidants can significantly lower the risk of several chronic diseases such as heart diseases, cancers and cataracts. The fungal pigments are of considerable interest in nutrition because of their role as antioxidants and potential for preventing or delaying degenerative diseases and for enhancing immune responses in animals and humans<sup>7</sup>. Recently, there is growing urge to discover natural antioxidants from microorganisms and microbial pigments are generally safer than the chemical and synthetic formulations. The pigments produced by *Penicillium* and *Monascus* (monascorubrine and monascorubramine) are structurally similar. Lovastatins or monacolins produced by *Penicillium*, *Monascus*, *Aspergillus* and *Rhizopus* inhibits cholesterol biosynthesis by binding to catalytic site of HMG-CoA reductase a key enzyme in cholesterol biosynthesis and scavenged DPPH radicals<sup>8,9,10,11</sup>. In the present study we report the antioxidant potential and characterization of fungal red pigment extracted from *P.purpurogenum*.

## MATERIALS AND METHODS

### Microorganism and culture conditions

The microorganism used in this study was isolated from soil, collected from Tiger reserve Parambikulam, Kerala, India. Stock cultures were maintained on potato dextrose agar (PDA) slants at 4°C after being incubated at 30°C for 5-7 days. PP-O production medium was used as fermentation medium to assess the pigment production. PP-O production medium containing 20 g of soluble starch and 2 g of yeast extract per liter of 50 mM citric acid/Na<sub>3</sub> citrate buffer with pH 5.0.

### Extraction and estimation of red pigment yield

Pure culture of *P. purpurogenum* from PDA slants was transferred into 250 ml Erlenmeyer flasks containing 100 ml of the growth medium.

After cultivation of 5-7 days, about 1 ml of the culture broth was dissolved with 5 ml of 90 % (v/v) methanol. The solvent and sample were kept on a rotary shaker at 200 rpm for 1 h, allowed to stand for 15 min and filtered through Whatman filter paper (47 mm). The clear supernatant was collected and after dilution, absorbance of red pigment was measured at 500 nm using Hitachi U-2000 spectrophotometer (M/s. Hitachi Ltd., Tokyo, Japan). The absorbance values were converted into pigment units using by the following formula<sup>12</sup>:

$$\text{Pigment yield (colour value units ml}^{-1}\text{)} = \frac{\text{OD} \times \text{Dilution} \times \text{Total volume of solvent}}{\text{Amount of sample (ml)}}$$

### Antioxidant activity

Lipid peroxidation, Hydroxy radical scavenging and nitric oxide scavenging antioxidant power assays were used to determine antioxidant activity of red pigment by *P. purpurogenum* that was carried out by following standard method<sup>13-15</sup>.

### Characteristics of red pigment

#### Thin layer chromatography (TLC)

Pigments were detected by thin-layer chromatography using silica gel plates and developed in n-butanol:acetic acid:water (12:30:50 v/v)) as a mobile phase and determined their R<sub>f</sub> values. The R<sub>f</sub> values is a mathematical representation of the ratio of the distance travelled by the solvent<sup>16</sup>. After TLC separation, the plates were kept in room temperature for drying. The spots in the TLC plates were detected using UV light.

#### Detection of chemical groups by Gas chromatography and mass spectrometry (GC-MS)

GC-MS analysis was carried out to identify the pigment compounds present in the TLC separated bands by using DB-5 column with a length of 30m (M/s. Perkin Elmer - Clarus 500). Helium used as a carrier gas with flow rate at 1 ml min<sup>-1</sup>.

#### Structure determination of pigment by Fourier Transformer Infra Red (FT-IR) Spectroscopy

To confirm the structure of fungal red pigment, FT-IR spectrometer (M/s. Impact 400D, Nicolet, Madison, WI) was used to measure the infrared spectra of extract solution in the wave

number of 400-4000  $\text{cm}^{-1}$  at room temperature. For each IR spectrometer samples 32 scans at 4  $\text{cm}^{-1}$  resolution was collected in the transmittance mode.

## RESULTS AND DISCUSSION

Antioxidant compounds in food play an important role as a health protecting factor. Scientific evidence suggests that antioxidants reduce the risk for chronic diseases including cancer and heart diseases. The main characteristic of an antioxidant is its ability to trap free radicals. Highly reactive free radicals and oxygen species are present in biological systems from a wide variety of sources. These free radicals might oxidize nucleic acids, proteins; lipid or DNA to initiate regenerative diseases. Estimation of Lipid peroxidation is the most commonly used method for determining antioxidant activity to measure the inhibition of linoleic acid. Scavenging of free radicals has been known as an established phenomenon in inhibition of lipid peroxidation which otherwise can be deleterious to cellular function. The antioxidant activities of the fungal pigment increased with increasing the concentration of the pigment extract of *P. purpurogenum* (Fig. 1). At 5  $\text{mg ml}^{-1}$ , *P. purpurogenum* exhibited 14.40 per cent of

antioxidant activity. However, antioxidant activity of standard BHA was 41.25, 43.37, 65.42 and 70.05 per cent at 5, 10, 15 and 20  $\text{mg ml}^{-1}$  respectively. These results are in accordance with the literature data of Tseng *et al.* (2006)<sup>17</sup>.

The hydroxyl radical is one of the representative reactive oxygen species generated in the body system. Hydroxyl radical produced may cause lipid peroxidation, sugar fragmentation, base loss and leakage of DNA strand<sup>18,19</sup>. The results of this assay revealed that the red pigment extract at the concentration of 20  $\text{mg ml}^{-1}$  were considerably more active. The pigment extract was found to possess the hydroxyl radical scavenging activity in a dose dependent manner. It exhibited a maximum of 74.92 per cent activity at 20  $\text{mg ml}^{-1}$  (Fig. 2). It is apparent from the present study that the *P. purpurogenum* and pigment not only scavenges off the free radical but also inhibits the generation of free radicals.

Nitric oxide (NO) is an important chemical mediator generated by endothelial cells, macrophages, neurons etc., and involved in the regulation of various physiological processes<sup>20</sup>. Excess concentration of NO is associated with several diseases<sup>21</sup>. Oxygen reacts with the excess nitric oxide to generate nitrite and peroxynitrite anions, which act as free radicals<sup>22</sup>. The nitric oxide




Fig. 1. Lipid peroxidation activity of red pigment




Fig. 2. Hydroxy radical scavenging ability of red pigment

Table 1. TLC of methanolic extracts of red pigment

Pigment fraction	Colour of the pigment fraction	Distance travelled (cm)	Rf value (cm)	Absorption maximum ( $\lambda$ max)
1	Pinkish red	10.3	0.79	500
2	Orange	11.5	0.88	420
3	Yellow	12.2	0.93	420

radical scavenging activity of the fungal pigment extract was less than that of BHA, which was used as standard for all the assays. At 20 mg ml<sup>-1</sup> concentration the pigment extract of *P. purpurogenum* exhibited higher activity of 43.23 per cent of inhibition. Whereas BHA standard recorded 81.58 per cent of inhibition (Fig. 3).

#### Identification of fungal pigments by TLC, GC-MS and FT-IR

In the present study, best separation of the red pigment compounds was achieved with the mobile phase consisting n-butanol:acetic acid:water (12:30:50). The pigment fractions of *P. purpurogenum* were separated in to 3 bands (pinkish red, orange and yellow) whose R<sub>f</sub> values and  $\epsilon_{\text{max}}$  were presented in Table 1. Close agreement was obtained between absorption maxima of these fractions and R<sub>f</sub> values. The results are in accordance with the earlier reports<sup>23, 24</sup>.




Fig. 3. Nitric oxide scavenging ability of red pigment

GC-MS studies were performed to identify the pigment compounds present in the TLC separated bands of methanolic extract of *P. purpurogenum* culture. The peak was obtained for red, orange and yellow of TLC separated bands from *P. purpurogenum* in GC-MS. The compound present in the red band have the retention time of 31.65 min. The given library match for this compound may be 3-(9a-methyl-3-octanyl-2,9,dioxo-2, 7,9,9a-dihydro-2H-furo[3,2-g]isochromen-6yl)-acrylic acid. It has the formula C<sub>23</sub>H<sub>26</sub>O<sub>7</sub> established from GC-MS (Fig. 4). For the orange and yellow bands in GC-MS, the given library match for these compounds may be 6-ethyl-5 hydroxy-2,3,7-trimethoxy naphthoquinone and 2-tert-butyl-4-isopropyl-5-methylphenol respectively (Fig. 5 and 6). The results are similar to who reported that *Penicillium* sp. AZ has been found to produce PP-O when cultured in a medium composed of




Fig. 4. GC-MS chromatogram of band 1 of crude extract of *P. purpurogenum*




Fig. 5. GC-MS chromatogram of band 2 of crude extract of *P. purpurogenum*




Fig. 6. GC-MS chromatogram of band 3 of crude extract of *P. purpurogenum*




Fig. 7. FT-IR spectrum of *P. purpurogenum* pigment extract

soluble starch and yeast extract at pH 7. They reported that presence of an azaphilone skeleton in PP-O pigment<sup>25</sup>. *Penicillium purpurogenum* has been found to produce Monascus pigment homologs in culture with a specific medium<sup>26</sup>. This strain does not produce citrinin (mycotoxin), making *P. purpurogenum* a potentially valuable commercial source of natural food colourant.

FT-IR spectroscopy has widely been used for the characterization and identification of fungi, bacteria and yeasts which are hydrophilic microorganisms and can easily be suspended in water for sample preparation<sup>27</sup>. In the present study, the infra red spectra of red pigment showed the presence of hydrogen bonded OH groups at 3247  $\text{cm}^{-1}$ . The carbonyl stretching vibration frequency of the pigment is in the region of 1716  $\text{cm}^{-1}$  for red pigment (Fig. 7). The observed stretching frequencies are however close to phenol and quinones. The aromatic C=C stretching observed at 1423 and 1558  $\text{cm}^{-1}$  for red pigment, Similar results were also reported by Dhale and Vijay-Raj (2009) in *Penicillium* sp<sup>28</sup>. The IR spectrum showing broad stretching at 3379  $\text{cm}^{-1}$  for hydroxyl group of phenolics. In the IR spectrum stretching frequency were also observed at 1643 and 1414  $\text{cm}^{-1}$  assignable to C=C and C-H of the aromatic ring respectively.

### CONCLUSION

In recent years, production of natural food colourants through microbial fermentation is an

extensive area of investigation, since they overcome concerns of unfavorable side effects by synthetic colours. In this study, under *in vitro* condition red pigment efficiently inhibits lipid peroxidation, scavenged free radicals and hydroxyl radical scavenging activity exhibited 30.15 and 74.92 per cent inhibition at the concentration of 20  $\mu\text{g ml}^{-1}$ . Identification of compounds in methanolic extract of red pigment from *P. purpurogenum* shown that the presence of phenols and quinones. This study reveals that fungal pigments are a better source of natural pigment and antioxidants than synthetic dyes, therefore, the investigations of the antioxidant activity of natural products have created new ways for drug development to reduce the usage of synthetic drugs.

### ACKNOWLEDGEMENT

The authors greatly acknowledge the Indian council of Agricultural Research (ICAR) for the financial support granted under the scheme "Application of microorganisms in Agriculture and allied sciences (AMAAS)" to carry out this study.

### REFERENCES

1. Dufosse, L. Microbial production of food grade pigments. *Food Technol. Biotechnol.*, 2006; **44**: 313-321.
2. Mapari, S.A.S., Nielsen, K.L., Larsen, T.O., Frisvad, J.C., Meyer, A.S., Thrane, U. Exploring fungal biodiversity for the production of water-soluble pigments as potential natural food colourants. *Curr. Opin. Biotechnol.*, 2005; **16**:



- 231–238.
3. Duran, N., Tixeira, M.F.S., De Conti, R., Esposito, E. Ecological-friendly pigments from fungi. *Crit. Rev. Food Sci Nutr.*, 2002; **42**: 53–66.
4. Mapari, S.A.S., Meyer, A.S., Thrane, U. Colorimetric characterization for comparative analysis of fungal pigments and natural food colorants. *J. Agric. Food Chem.*, 2006; **54**(19): 7028–7035.
5. Halliwell, B., Gutteridge, M.C. Oxygen toxicity, oxygen radical, transition metals and disease. *Biochemical Journal*, 1984; **219**: 1–14.
6. Vaya, J., Aviram, M. Nutritional antioxidants: Mechanism of action, analyses of activities and medical applications. *Curr. Med. Chem- Imm. Endoc. and Metab. Agents*, 2001; **1**: 99–117.
7. Kirakosyan, A., Seymour, E., Kaufman, O.B., Warber, S., Bolling, S., Chang, S.C. Antioxidant capacity of polyphenolic extracts from leaves of *Crataegus laevigata* and *Crataegus monogyna* (Hawthorn) subjected to drought and cold stress. *J. Agric. Food Chem.*, 2003; **51**: 3973–3976.
8. Endo, A., Kuroda, M., Tsujita, Y. ML-236A, ML-236B, ML-236C, new inhibitors of cholestrogenesis produced by *Penicillium citrinum*. *Journal of Antibiotics (Tokyo)*, 1979; **29**: 1346–1348.
9. Albert, A.W. Lovastatin and Simvastatin-inhibitors of HMG-CoA reductase and cholesterol biosynthesis. *Cardiology*, 1990; **77**: 14–21.
10. Dhale, M.A., Divakar, S., Umesh-Kumar, S., Vijayalakshmi, G. Isolation and characterization of dihydromonacolin-MV from *Monascus purpureus* for antioxidant properties. *Applied Microbiology and Biotechnology*, 2007a; **73**: 1197–1202.
11. Dhale, M.A., Divakar, S., Umesh-Kumar, S., Vijayalakshmi, G. Characterization of dehydromonacolin-MV2 from *Monascus purpureus* mutant. *Journal of Applied Microbiology*, 2007b; **103**: 2168–2173.
12. Johns, M.R., Stuart, D.M. Production of pigments by *Monascus purpureus* in solid culture. *J. Ind. Microbiol.*, 1991; **8**(1): 23–28.
13. Lingnert, H., Vallentin, K., Eriksson, C.E. Measurement of antioxidative effect in model system. *J. Food Process. Preserv.*, 1979; **3**: 87–103.
14. Lloyd, R. V., Hanna, P. M., Mason, R.P. The origin of the hydroxyl radical oxygen in the Fenton reaction. *Free Radical Biol. Med.*, 1997; **22**(5): 885–888.
15. Marcocci, L., Maguire, J.J., Droy-Lefaix, M.T., Packer, L. The nitric oxide scavenging property of *Ginkgo biloba* extract EGb 761. *Biochem. Biophys. Res. Commun.*, 1994; **201**: 748–755.
16. Blanc, P.J., Loret, M.O., Sanberre, A.L., Pareilleuz, A., Prome, D., Laussac, J.P., Goma, G. Pigments of *Monascus*. *J. Food Sci.*, 1994; **59**: 862–865.
17. Tseng, Y.H., Yang, J.H., Chang, H.L., Lee, Y.L., Mau, J.L. Antioxidant properties of methanolic extracts from *Monascus* adlay. *Food chem.*, 2006; **97**: 375–381.
18. Xu, Q., Tao, W., Ao, Z. Antioxidant activity of vinegar melanoidins. *Food Chem.*, 2007; **102**: 841–849.
19. Morales, F.J., Jimenez-Perez, S. Peroxyl radical scavenging activity of melanoidins in aqueous systems. *Eur. Food Res. Technol.*, 2004; **218**(6): 515–520.
20. Lata, H. Ahuja, G.K. Role of free radicals in health and disease. *Ind. J. Physiol. Allied Sci.*, 2003; **57**: 124–128.
21. Ialenti, A., Moncada, S., Di Rosa, M. Modulation of adjuvant arthritis by endogenous nitric oxide. *Br. J. Pharmacol.*, 1993; **110**: 701–705.
22. Sainani, G.S., Manika, J.S., Sainani, R.G. Oxidative stress- a key factor in pathogenesis of chronic diseases. *Med. Update*, 1997; **1**: 1–5.
23. Ogihara, J., Oishi, K. Effect of ammonium nitrate on the production of PP-V and monascorubrin homologues by *Penicillium* sp. AZ. *J. Biosci. Bioeng.*, 2002; **93**: 54–59.
24. Arai, T., Koganei, K., Umemura, S.R., Kojima, Kato, J., Kasumi, T., Ogihara, J. Importance of the ammonia assimilation by *Penicillium purpurogenum* in amino derivative *Monascus* pigment, PP-V, production. *AMB Express*, 2013; **3**(19): 1–7.
25. Ogihara, J., Kato, J., Oishi, K., Fujimoto, Y., Eguchi, T. Production and structural analysis of PP-V, a homologue of monascorubramine, produced by a new isolate of *Penicillium* sp. *J. Biosci. Bioeng.*, 2000a; **90**: 549–554.
26. Ogihara, J., Kato, J., Oishi, K., Fujimoto, Y. Biosynthesis of PP-V, a monascorubramine homologue by *Penicillium* sp. AZ. *J. Biosci. Bioeng.*, 2000b; **90**: 678–680.
27. Fischer, G., Braun, S., Thissen, R., Dott, W. FT-IR spectroscopy as a tool for rapid identification and intra-species characterization of airborne filamentous fungi. *J. Microbiol. Methods*, 2006; **64**: 63–77.
28. Dhale, M.A., Vijay raj, A.S. Pigment and amylase production in *Penicillium* sp. NIOM-02 and its radical scavenging activity. *Int. J. Food Sci. Technol.*, 2009; **44**(12): 2424–2430.