

## Management of Pomegranate Wilt Complex Caused by *Ceratocystis fimbriata* and *Meloidogyne incognita*

Shreeshail Sonyal<sup>1</sup>, V.B. Nargund<sup>2</sup>, Yallappa Jagarkal<sup>3</sup>, V.I. Benagi<sup>2</sup>, K.B. Palanna<sup>1</sup>,  
Madhu S. Giri<sup>2</sup>, Anil Pappachan<sup>1</sup>, H. Shivalingappa<sup>1</sup>, H.S. Mahesha<sup>1</sup>,  
Devanshu Dev<sup>1</sup> and M.E. Puneeth<sup>1</sup>

<sup>1</sup>Department of Plant Pathology, UAS, GKVK, Bangalore, India.

<sup>2</sup>Department of Plant Pathology, UAS, Dharwad, India.

<sup>3</sup>Department of Agricultural Microbiology, UAS, GKVK, Bangalore, India.

<https://doi.org/10.22207/JPAM.10.1.19>

(Received: 11 October 2015; accepted: 24 December 2015)

In this experiment 45<sup>th</sup> day was crucial day for wilting. Branches were rapidly wilted in all treatment. After 45<sup>th</sup> day gradual yellowing and wilting was noticed very slowly due to effect of chemicals, bio agents and plant product, micronutrient insecticide. Among the chemicals propiconazole was more effective followed by Difenconazole. However in bio agents *T. harzianum* and *P. fluorescens* were effective than alone. The propiconazole and difenconazole was significantly superior than remaining treatments after 120<sup>th</sup> day propiconazole (0.00%), difenconazole (0.00%) and *T. harzianum* + *P. fluorescens* (0.00%) are on par with each other. The percent survival of plants in captan+ carbendazim (83.34%), *T. harzianum* (83.34%), Copper oxy chloride (83.34%) and mancozeb (83.34%) were on par with each other. Followed by neemcake (66.66%) and vitavax power (66.66%). In control (83.33%) totally showing least percent of survival of plants.

**Key words:** 45<sup>th</sup> day, 120<sup>th</sup> day, Propiconazole, wilt and crucial day.

Pomegranate (*Punica granatum* L.) is an ancient fruit, belonging to the smallest botanical family puniceae. Pomegranate is a native of Iran, where it was first cultivated in about 2000 BC. Sweet varieties are mildly laxative, sour types are good for curing inflammation of stomach and heartache. In India, there is a common adage 'EkAnar Sau Bimar' meaning one fruit cures hundred diseases. Area under pomegranate is increasing worldwide because of its hardy nature, wider adaptability, and drought tolerance, higher yield levels with excellent keeping quality and remunerative prices in domestic as well as export market. It thrives well in dry tropics and sub-tropics

and comes up very well in soils of low fertility status as well as on saline soil.

In India, it is regarded as a "vital cash crop", grown in an area of 1, 16,000 ha with a production of 89,000 MT with an average productivity of 7.3 MT (Srivastava and Umesh, 2008). Among the different states growing pomegranate, Maharashtra is the largest producer occupying 2/3<sup>rd</sup> of total area in the country followed by Karnataka, Andhra Pradesh, Gujarat and Rajasthan. Karnataka state has the distribution of cultivating pomegranate under tropical condition in an area of 12,042 ha with a production of 1, 29, 547 tonnes. Where this crop has spread across different districts viz., Bijapur, Bellary, Bagalkot, Koppal, Chitradurga, Belgaum, Davangere, Tumkur, Bangalore and Gulbarga.

\* To whom all correspondence should be addressed.  
E-mail : shailgkvk2012@gmail.com

## MATERIAL AND METHODS

A field experiment was conducted during January to May 2010 at farmer field, Bandi under irrigated condition to find out the suitable fungicide for the management of wilt. The experiment was laid out in completely randomized block design (RCBD) with 12 treatments and replicated twice. In each treatment six plants were tagged and the efficacy of ten fungicides (four systemic, three non-systemic fungicides and two bio agents, one plant product and one insecticide, one micro nutrient) was tested with one untreated control. The fungicide solutions were prepared by dissolving known quantity of fungicide in water and eight litres per tree drenched. First drenching was taken 3<sup>rd</sup> January and second drenching with 15 days interval.

### Design of the layout

The experiment was laid out in a completely randomized block design with two replications. The treatments were randomly allotted to plots. The details are as follows.

Treatments: Twelve

#### The treatments are as follows

S. No.	Treatments	Common name	Conc. (%)
1	T1	Control	-
2	T2	Carbendazim	0.2
3	T3	Vitavax power	0.2
4	T4	Captan+ Carbendazim	0.2+0.2
5	T5	Neem cake	1kg
6	T6	<i>P. flourescens</i> + <i>T. harzianum</i>	10 <sup>6</sup> +10 <sup>6</sup>
7	T7	<i>T. harzianum</i>	10 <sup>6</sup>
8	T8	Mancozeb	0.3
9	T9	Copper oxy chloride	0.3
10	T10	Difenoconazole	0.2
11	T11	Hexaconazole	0.2
12	T12	Propiconazole	0.2
1	Borax	32ml/tree	
2	Chlorpyriphos	64g /tree	

Number of plants tagged/treatment: Six

Spacing : 4.25 m x 3.65m

Variety : Kesar

The area under disease progression curve (AUDPC) was calculated using the formula of Nagarajan and Muralidharan (1995).

$$\text{AUDPC} = S \frac{1}{2} (Y_1 + Y_{i-1}) \times d$$

J PURE APPL MICROBIO, 10(1), MARCH 2016.

Y<sub>i</sub> = Disease incidence at i<sup>th</sup> day or evaluation

K = Number of successive evaluation

D = Interval between i and i-1 evaluation of disease.

## RESULTS

The results of the field experiment conducted in a farmer's field at Bandi village, Yalburga taluk (Koppal District) are presented in Table 1.



In this experiment 45<sup>th</sup> day was crucial day for wilting (Plate 1). Branches were rapidly wilted in all treatment. After 45<sup>th</sup> day gradual yellowing and wilting was noticed very slowly due to effect of chemicals, bio agents and plant product, micronutrient insecticide. Among the chemicals propiconazole was more effective followed by Difenoconazole. However in bio agents *T. harzianum* and *P. flourescens* were effective than alone.

From the Table 1 (Fig. 1) results indicates that propiconazole was significantly superior than remaining treatments. Propiconazole treated plants were recovered at 105<sup>th</sup> day, followed by difenoconazole the plants were recovered at 120<sup>th</sup> day. These two chemicals were effective than remaining treatments. Carbendazim (57.89%), vitavax powers (58.82%) are on par with each other and showed least efficacy. Captan + carbendazim (44.45%), *T. harzianum* (44.11%) Copper oxychloride (44.44%) are on par with each other. Followed by neemcake (52.94%) and *T. harzianum* + *P. flourescence* (48.48%) are on par with each other. The control showing maximum percent of wilting (97.61%). Before drenching all plants showing partial wilting and yellowing but none of the plants completely wilted in all treatments. After 120<sup>th</sup> day propiconazole (0.00%), difenoconazole (0.00%) and *T. harzianum* + *P. flourescence* (0.00%) are on par with each other. Captan + carbendazim (16.66%), *T. harzianum* (16.66%), Copper oxychloride (16.66%) and mancozeb (16.66%) were on par with each other. Followed by neemcake (33.34%) and vitavax power (33.34%). In control (83.33%) was showing highest percent of wilting.

The propiconazole and difenoconazole was significantly superior than remaining treatments after 120<sup>th</sup> day propiconazole (0.00%), difenoconazole (0.00%) and *T. harzianum* + *P. flourescence* (0.00%) are on par with each

**Table 1.** Per cent survival of branches and plants after 120 days

S. No.	Treatments	Per cent wilting of branches before drenching	Per cent wilting of branches after 120 days	Per cent survival	Per cent wilting of plants before drenching	Per cent wilting of plants after 120 days	Per cent survival
1	Control	14.28	97.61	2.39	0.00	83.33	16.67
2	Carbendazim	10.52	57.89	42.11	0.00	50.00	50.00
3	Vitavax power	11.76	58.82	41.18	0.00	33.34	66.66
4	Captan + Carbendazim	11.12	44.45	55.55	0.00	16.66	83.34
5	Neem cake	8.82	52.94	47.06	0.00	33.34	66.66
6	<i>T. harzianum</i> + <i>P. fluorescens</i>	12.12	48.48	51.52	0.00	0.00	100
7	<i>T. harzianum</i>	11.76	44.11	55.89	0.00	16.66	83.34
8	Mancozeb	11.76	41.17	58.83	0.00	16.66	83.34
9	Copper oxy chloride	11.12	44.44	55.56	0.00	16.66	83.34
10	Difenconazole	9.38	0.00	100	0.00	0.00	100
11	Hexaconazole	10.52	42.10	57.9	0.00	16.66	83.34
12	Propiconazole	12.50	0.00	100	0.00	0.00	100

**Fig. 1.** Percent survival of branches after 120 days**Fig. 2.** Percent survival of plants after 120 days**Plate 1.** General view of experimental plot

other. The percent survival of plants in captan+carbendazim (83.34%), *T. harzianum* (83.34%), Copper oxy chloride (83.34%) and mancozeb (83.34%) were on par with each other. Followed by neem cake (66.66%) and vitavax power (66.66%). In control (83.33%) totally showing least percent of survival of plants.

## DISCUSSION

Protection of crop plants from disease causing agents has been the focal point of the scientists in dealing with plant pathogens. In-depth analysis and realization about the startling features of microbial ecology have impelled workers to replace term control with management. No single method of pest control has given a lasting solution. Prohibitive costs of chemicals and their adverse ecological impact a major compulsion for

diversions of research priorities from chemical methods to other alternatives. The successive shifting of priorities from chemical to cultural and currently integrated management demonstrates the elasticity of scientific ideas (Khan and Parvatha Reddy, 1993).

Present study clearly indicated that the - propiconazole was significantly superior to all other treatments followed by difenconazole and *T. harzianum* + *P. fluorescens*, because out of six plants six plants were completely green. In propiconazole plant recovery was faster than other treatments. In *T. harzianum* one plant was wilted and four were remain as green. In carbendazim + captan only one plant was wilted two plants remain as green out of six plants on par with mancozeb, Copper oxy chloride and hexaconazole. In vitavax power two plants were died and three plants remain as green on par with Neem cake. In carbendazim three plants were died and single plant remain as green. Controls Out of six plants 5 plants were died and only one branch remains as yellow, 1<sup>st</sup> plant wilted at 45<sup>th</sup> day. The present investigations are confirmation of findings, Jadhav and Sharma (2009) who has reported that drenching of diseased plants and also adjacent healthy plants with carbendazim (0.2 %), propiconazole (0.2 %) + chlorpyrifos (0.2 %) is effective method of disease management.

The present findings were confirmed by Sharma and Mehrotra (1984) observed that the use of carbendazim and benomyl gave good control of *C. paradoxa*. The budded sugarcane setts treated with fungicides such as agalol, carbendazim, tridimefon or benomyl, good growth of sett with minimum sett rot incidence (Taylor and Ryam, 1984). Two fungicides topsin and busan were

superior against *C. paradoxa* than emisan (Pandu *et al.*, 1986). The effective control of sett rot of sugarcane with carbendazim (Rehman *et al.*, 1989).

## REFERENCES

1. Jadhav, V. T. and Sharma, K. K., Integrated management of disease in pomegranate. Paper Presented In: 2nd *Inter. Symp. Pomegranate and minor including Mediterranean Fruits*, Univ. Agric. Sic., Dharwad, 2009; 23-27, 48-52 pp.
2. Khan, R. M. and Parvatha Reddy, P., Management of disease complex. In *Nematode Interactions*, Ed. Khan, M.N., Chapman and Hall, London, 1993; 345-365.
3. Nagarajan, S and Muralidharan, K., *Dynamics of Plant Diseases*. Allied publishers Limited, New Delhi, India, 1995; 2.
4. Pandu, S. R., Rao, S. V., Sharma, M. N., and Subbaiyya, J., Effectiveness of non-mercurial fungicides for the control of pineapple disease of sugarcane. *Indian Phytopath.* 1986; **39**: 306-308.
5. Rehman, S., Kabir, A. K. M., Talukder, I., and Iqbal, M., Efficacy of sett treating fungicides in sugarcane production in Bangladesh, *Bangla. J. Sug.* 1989; **11**: 72-77.
6. Sharma, S. K., and Mehrotra, R.S., Chemical controls of stem rot disease of rice caused *Sclerotium oryzae*. *Indian Phytopathol.*, 1984; **37**: 390.
7. Srivastava and Umesh, C., Horticulture in India: a success story. *Crop Care*, 2008; **34**: 23-32.
8. Taylor, P.W.J., and Ryam, C.C., Propiconazole fungicide as a sett treatment for the control of pineapple disease. *Sugarcane*, 1984; **5**: 5-8.

© The Author(s) 2016. Open Access. This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.