

Phylogenetic Study of *Escherichia coli* Isolated from Clinical Samples in Hilla City, Iraq

Noor S. K. Al-Khafaji* and Azhar A. L. Al-Thahab

University of Babylon, College of Sciences-Biology Dept. Hilla, Iraq.

<http://dx.doi.org/10.22207/JPAM.11.4.16>

(Received: 26 September 2017; accepted: 10 November 2017)

Escherichia coli clones, designated as extraintestinal pathogenic *E. coli* (ExPEC), are responsible for extraintestinal infections. Phylogenetic analysis has shown that *E. coli* is composed of four main phylogenetic groups (A, B1, B2 and D) and six subgroups (i.e. A0, A1, B2₂, B2₃, D1 and D2). Group A and B1 are generally associated with commensals, whereas group B2,D is associated with extra-intestinal pathotypes. In the present study, a total of 53 *E. coli* isolates, isolated from human clinical samples, were used. Phylogenetic grouping was done based on the PCR method using primers targeted at three genetic markers, *chuA*, *yjaA* and *TspE4.C2*. According to PCR-based phylotyping, subgroup B2₃ contained the majority of the collected isolates (40 isolates, 75.47 %), followed by subgroups A1 and B2₂ (4 isolates for each subgroup, 7.55 %), followed by subgroups D2 (3 isolates, 5.66 %) and A0, D1 (1 isolate for each subgroup, 1.88 %). No isolates were found to belong to group B1. Based on the results, the majority of isolates were extra-intestinal pathotypes. Therefore, the role of *E. coli* in human infections including urinary tract infections, septicemia, vaginitis and meningitis should be considered for further research.

Keywords: *Escherichia coli*, Clinical Samples, Iraq.

Extraintestinal pathogenic *E. coli* (ExPEC) represents a distinct group of pathogenic *E. coli* that causes most of the extraintestinal *E. coli* infections (urinary tract infection, prostatitis, bacteremia, septicemia, and neonatal meningitis, vaginosis) human infections (Donnenberg *et al.*, 2002; Gordon and Cowling, 2003; Al-Khaqani *et al.*, 2016; Abdulla *et al.*, 2016). ExPEC isolates are genetically distinct from commensal *E. coli* found in the intestinal flora. Phylogenetic analyses have revealed that *E. coli* isolates are composed of four main phylogenetic groups (A, B1, B2, and D) (Gordon, 2004; Al-Khaqani *et al.*, 2016). Isolates of each of the four groups have different phenotypic features, causing their ability to exploit different sugars, antibiotic-resistance profiles and growth

rate-temperature relationships. The distribution (presence/absence) of a variety of genes thought to enable a strain to cause extra-intestinal disease also varies among isolates of the four phylo-groups (Johnson *et al.*, 2001). Several studies have shown the relation between phylogeny and pathogenicity of *E. coli* isolates (Bashir *et al.*, 2012; Escobar-Paramo *et al.*, 2004). Bearing in mind that most commensal isolates belong to A and B1 groups (Duriez *et al.*, 2001), Phylogenetically and epidemiologically ExPEC are potentially different from those of intestinal pathogenic and commensal isolates (Smith *et al.*, 2007). Most of the ExPEC isolates phylogenetically belong to B2 and to a lesser extent D groups and are equipped with various virulence factors that help these isolates during different mode of infection mechanisms like adhesion, invasion of host tissues, escape host defence mechanisms, signaling and production of different toxins interfering host cellular functions

* To whom all correspondence should be addressed.
E-mail: noorvaccine@gmail.com

thereby promoting extraintestinal infection in both normal and immune compromised hosts (Dobrindt and Hacker, 2008; Wiles *et al.*, 2008; Al-Dahmoshi *et al.*, 2016)

The aim of this study was to investigate the phylo-genetic groups of *E. coli* isolated from different clinical samples which includes urine, vaginal , seminal fluid and wound swap in Hilla, Iraq using a molecular primer.

MATERIAL AND METHODS

A total of 200 various clinical samples represent by 50 urine samples from patients with urinary tract infection, 50 high vaginal swap from pregnant and non-pregnant women suffering from vaginosis, 50 seminal fluid samples of male suffering from bacteriospermia and 50 wound swabs samples . All samples were obtained from patients or individuals who were admitted to Babylon Hospital for Maternal and Pediatrics, and to Al-Hilla Surgical Teaching Hospital in Babylon city (Iraq) during the period from May to August 2016. In order to isolate *E. coli*, samples were directly inoculated on MacConkey agar (Himedia/India) plates. After overnight incubation at 37°C, lactose fermenting colonies, Gram-negative, oxidase negative bacilli transferred to UTI Chromogenic medium (Condalab/Spain) and Eosin methylene blue agar (Himedia/India) to confirm *E. coli* isolates.

DNA extraction for gram negative bacteria

Typical *E. coli* colonies (with metallic green color on Eosin methylene blue agar and pink color on UTI Chromogenic medium) were

grown in LB broth(Condalab/Spain) at 37°C for 18 h, and following the protocols of Favor Prep Genomic DNA Mini Kit (Blood/Cultured Cell) (Favorgen/Taiwan). The extracted DNA checked using Agarose gel electrophoresis (0.7% in TBE buffer) (Condalab/Spain) and then visualized using and gel documentation (Vilber/France).

Detection of phylogeny groups by PCR

PCR was conducted to determine the phylogenetic grouping of the isolates by targeting three genes, *chuA*, *yjaA* and *TspE4.C2* using 20iL reaction mix (IntronBio/Korea) (Clermont *et al.*, 2000). Thermal cycler conditions were as follows: 95°C for 4 min, 30 cycles of (denaturation at 94°C for 30sec.), (annealing at 59 °C for 30sec.), (extension at 72°C for 30sec). and final extension at 72°C for 5 min. Agarose gel electrophoresis (1.5% in TBE buffer) and gel documentation (Vilber/France) were used to visualized and document the PCR products. The amplicon sizes were 279 bp for *chuA*, 211 bp for *yjaA* and 152 bp for *TspE4C2* were recorder using 100bp ladder (IntronBio/Korea). After electrophoresis the gel was photographed under UV light. The results allowed the classification of isolates into either one of the four major phylogroups (A, B1, B2, or D) (Abdallah *et al.*, 2011; Gordon *et al.*, .2008).

RESULTS AND DISCUSSION

This study was carried out to expose the phylogeny of *E. coli* isolated from different clinical samples to investigate the source of these isolates whether they are intestinal or extraintestinal. However, the phylogenetic groups of *E. coli*




Fig. 1. 1.5% Agarose gel electrophoresis at 72 volt for 60 minutes of PCR to *chuA* amplicon (279bp); lane M represent DNA marker size(100bp). Lane(1-9) represent some of positive *E. coli* isolates.

isolated from clinical samples were detected by identifying the presence of specific PCR amplified fragments (*chuA*, *yjaA*, and *TspE4.C2*) (Figures 1,2,3) respectively.

A total of 53 *E. coli* isolates represented by (12 vaginal, 17 UTI, 10 seminal fluid and 14 wound swap) samples and according to the presence or absence of these genes were assigned to one of four phylogenetic groups (i.e. A, B1, B2 and D) group A and B1 (intestinal groups); B2 and D (extraintestinal groups) and six subgroups (i.e. A₀, A₁, B_{2,2}, B_{2,3}, D₁ and D₂). According to PCR-based phylotyping the result showed that all vaginal samples, 14 urine samples, 8 of wound samples and 6 of seminal fluid samples were belong to B_{2,3} subgroup, 1 of urine samples and 3 of wound samples were belong to B_{2,2} subgroup, 2 of urine samples and 1 of seminal fluid samples were belong to D₂ subgroups, 2 of wound samples and 2 of seminal fluid samples were belong to A₁

subgroup, 1 of wound swap samples was belong to A₀ subgroup and 1 of seminal fluid samples was belong to D₁ subgroup. Also the results showed that the subgroup B_{2,3} contained the majority of the collected isolates (40 isolates, 75.47%), followed by subgroups A₁, B_{2,2} (4 isolates for each subgroup, 7.55%) then subgroup D₂ (3 isolates, 5.66%) and A₀, D₁ (1 isolate for each subgroup, 1.88%). No isolates were found to belong to group B1 (Table 1 and 2)

DISCUSSION

The niche of commensal *E. coli* is the mucous layer of the colon. However, there are *E. coli* clones that are distinct from the intestinal commensal *E. coli*, possessing specific fitness and virulence attributes which allow adaptation to other niches (e.g. urinary tract, central nervous system, blood) and confer enhanced ability to cause a broad




Fig. 2. 1.5% Agarose gel electrophoresis at 72 volt for 60 minutes of PCR to *yjaA* amplicon (211bp); lane M represent DNA marker size(100bp). Lane(1-9) represent some of positive *E.coli* isolates




Fig. 3. 1.5% Agarose gel electrophoresis at 72 volt for 60 minutes of PCR to *TspE4.C2* amplicon (152bp); lane M represent DNA marker size(100bp). Lane(1-9) represent some of positive *E.coli* isolates

Table 1. Percentage for isolation of extraintestinal *E. coli*

Sample	Urine	Vaginal Swab	Seminal fluid	Wound Swab	Total
No. of Samples	50	50	50	50	200
<i>E. coli</i> No.(%)	17(34%)	12(24%)	10(20%)	14(28%)	53(26.5%)
ExPEC, No.(%)	17(34%)	12(24%)	8(16%)	11(22%)	48 (24%)

Table 2. Percentage of *E. coli* isolates for each phylogenetic subgroups

Phylogenetic groups N(%)	Phylogenetic subgroups	N (%)	Total (%)	
Intestinal Groups	Group A	Subgroup A0chuA - / yjaA - / TspE4.C2 -	1 (1.88)	5 (9.43)
		Subgroup A1chuA - / yjaA + / TspE4.C2 -	4 (7.55)	
Extraintestinal Groups	Group B1	chuA - / yjaA - / TspE4.C2 +	0 (0)	48 (90.57)
	Group B2	Subgroup B22chuA + / yjaA + / TspE4.C2 -	4 (7.55)	
		Subgroup B23chuA + / yjaA + / TspE4.C2 +	40 (75.47)	
	Group D	Subgroup D1chuA + / yjaA - / TspE4.C2 -	1 (1.88)	
	Subgroup D2chuA + / yjaA - / TspE4.C2 +	3 (5.66)		
Total (%)		53 (100)		

spectrum of disease in extraintestinal sites (Johnson and Russo,2005) . ExPEC isolates usually belong to phylogenetic group B2 and to a lesser extent to group D, whilst commensal isolates are derived from groups A and B1 (Escobar-Páramo et al., 2004). Therefore, the rapid PCR-based phylogenetic typing developed by Clermont et al.(Clermont et al .,2000) has proven useful for rapidly screening ExPEC. According to this method, about more than two-thirds of the analyzed isolates in this study belonged to group B2 (83.01%) and this is larger than the studies of Obata-Yasuoka et al, 2002 (76%) ,Watt et al,2003 (68%) and Hilbert et al, 2008 (62%). A Compared to the data reported by these reserchers, group D isolates had a lower prevalence in this study (7.54% vs. 16%, 16%, and22%, respectively) , but the commensal phylogenetic group A exhibited nearest prevalence among the isolates investigated by this study (9.43% vs. 8%, 12% and 8% respectively). These differences in distribution of the phylogenetic groups among the isolates of geographically distinct populations in different studies may be due to the health status of the host, geographic climatic conditions, dietary factors, the use of antibiotics, or host genetic factors. (Duriez et al., 2001) . Based on the results of this study , the majority of isolates were

extraintestinal pathotypes. Therefore, the role of *E. coli* in human infections including urinary tract infections, septicemia, vaginitis and meningitis should be considered for further researches.

REFERENCES

1. Abdallah KS, Cao Y, Wei DJ. Epidemiologic investigation of extraintestinal pathogenic *E. coli* (ExPEC) based on PCR phylogenetic group and fimH single nucleotide polymorphisms (SNPs) in China. *Int J Mol Epidemiol Genet*, 2011; **2**: 339-353.
2. Abdulla, A.A., Al-Dahmoshi, H.O.M., Abed, T.A. and Muttaleb, W.H. Characterization of multidrug resistant carbapenemases-producing *Escherichia coli* and *Klebsiella pneumoniae* isolates from urinary tract infection. *Journal of Chemical and Pharmaceutical Sciences*, 2016; **9**(3):1116-1120.
3. Al-Dahmoshi, H.O.M., Almamoori, A.M.J. and Al-Khafaji, N.S.K. Investigation of CusCFBA pump among uropathogenic *Escherichia coli* (UPEC) Isolated from women with cystitis, Iraq. *Asian Journal of Pharmaceutical and clinical research*, 2016; **10**(1).
4. Al-Khaqani, M.M., Alwash, M.S. and Al-Dahmoshi, H.O. Phylogrouping, antibiotics susceptibility and biofilm formation among

- cervico-vaginal *Escherichia coli* (CVEC) isolated for female in Hilla City, Iraq. *Malaysian Journal of Microbiology*, 2016; **12**(4).
5. Bashir S, Haque A, Sarwar Y, Anwar A, Anwar M. Virulence profile of different phylogenetic groups of locally isolated community acquired uropathogenic *E. coli* from Faisalabad region of Pakistan. *Ann Clin Microbiol Antimicrob.* 2012; **11**:23.
 6. Clermont O, Bonacorsi S, Bingen E. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl. Environ. Microbiol.*, 2000; **66**: 4555-4558.
 7. Dobrindt U. and Hacker J. Targeting virulence traits: potential strategies to combat extraintestinal pathogenic *E. coli* infections. *Current opinion in microbiology*, 2008; **11**: 409-413.
 8. Donnenberg M. *Escherichia coli* virulence mechanisms of versatile pathogen. Elsevier Science, San Diego Calif. : Academic Press; 2002. pp. xxi-xxv.
 9. Duriez P, Clermont O, Bonacorsi S, Bingen E, Chaventre A, Elion J, Picard B, Denamur E. Commensal *Escherichia coli* isolates are phylogenetically distributed among geographically distinct human populations. *Microbiology*, 2001; **147**: 1671-1676.
 10. Escobar-Paramo P, Clermont O, Blanc-Potard A, Bui H, Le Bouguenec C, Denamur E. A specific genetic background is required for acquisition and expression of virulence factors in *Escherichia coli*. *Mol Biol Evol.*, 2004; **21**: 1085-1094.
 11. Gordon D, Cowling A. The distribution and genetic structure of *Escherichia coli* in Australian vertebrates: host and geographic effects. *Microbiology*, 2003; **149**: 3575-3586
 12. Gordon DM, Clermont O, Tolleyand H, Denamur E. Assigning *Escherichia coli* isolates to phylogenetic groups: multi-locus sequence typing versus the PCR triplex method. *Environ Microbiol*, 2008; **10**: 2484-2496.
 13. Gordon DM. The influence of ecological factors on the distribution and genetic structure of *Escherichia coli*. In *Escherichia coli and Salmonella typhimurium*. American Society for Microbiology 2004; [[http:// www.ecosal.org/ecosal/index.jsp](http://www.ecosal.org/ecosal/index.jsp)].
 14. Hilbert DW, Paulish TE, Mordechai E, Adelson ME, Trama JP. O serogroups, phylogeny, and virulence factors of cervicovaginal and rectal *Escherichia coli* isolates *Eur. J Clin Microbiol Infect Dis*, 2008; **27**(12):1265-8.
 15. Johnson JR, Delavari P, Kuskowski M, Stell AL. Phylogenetic distribution of extraintestinal virulence-associated traits in *Escherichia coli*. *J Infect Dis*, 2001; **183**:78-88.
 16. Johnson JR, Russo TA. Molecular epidemiology of extraintestinal pathogenic (uropathogenic) *Escherichia coli*. *Int J Med Microbiol.*, 2005; ; **295**(6-7):383-404
 17. Obata-Yasuoka M, Ba-Thein W, Tsukamoto T, Yoshikawa H, Hayashi H. Vaginal *Escherichia coli* share common virulence factor profiles, serotypes and phylogeny with other extraintestinal *E. coli*. *Microbiology*, 2002; **148**(Pt 9):2745-52.
 18. Orskov F, Orskov I. *Escherichia coli* serotyping and disease in man and animals. *Can J Microbiol*, 1992; **38**: 699-704.
 19. Smith J.L., Fratamico P.M. and Gunther N.W. Extraintestinal pathogenic *Escherichia coli*. *Foodborne pathogens and disease*, 2007; **4**: 134-163.
 20. Watt S, Lanotte P, Mereghetti L, Moulin-Schouleur M, Picard B, Quentin R. *Escherichia coli* isolates from pregnant women and neonates: intraspecies genetic distribution and prevalence of virulence factors. *J Clin Microbiol.*, 2003; **41**(5):1929-35.
 21. Wiles T.J., Kulesus R.R. and Mulvey M.A. Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Experimental and molecular pathology*, 2008; **85**: 11-19.