

Molecular Characterization and Evaluation of Two Potential Mosquitocidal *Lysinibacillus* Strains from Himalayan Valley Kashmir

Reyaz Ahmad Lone and P. Indra Arulselvi*

Plant and Microbial Biotechnology Laboratory,
Department of Biotechnology Periyar University, India.

<http://dx.doi.org/10.22207/JPAM.11.4.20>

(Received: 02 October 2017; accepted: 20 November 2017)

In an exertion to isolate natural non-harmful mosquitocidal bacteria, 158 samples of soil were collected from various habitats of Himalayan valley Kashmir. A total of 450 bacteria were screened for mosquitocidal activities against three epidemiological disease causing vectors: *Aedes aegypti*, *Culex quinquefasciatus* and *Anopheles stephensi* larvae/pupae. Out of 450 bacteria screened, none had shown pupicidal activity. However, two isolates KS2-15 and KS2-13 exhibited mosquito larvicidal activity against *C. quinquefasciatus* (LC50: 1.36×10^3 spores/mL; 1.41×10^3 spores/mL respectively) and *A. stephensi* (LC50: 2.14×10^3 spores/mL; 2.11×10^3 spores/mL correspondingly). These two isolates were identified, morphologically, biochemically and comparative investigation of 16S rRNA gene sequences, as *Lysinibacillus sphaericus* (previously *Bacillus sphaericus*). Sodium dodecyl sulfate polyacrylamide gel electrophoresis analysis of spore crystal mixture of each strain revealed two major bands of around 51.4 kDa and 41.9 kDa. PCR of mosquitocidal toxin genes showed the presence of *binA* and *binB* genes in both the strains. Comparative amino acid sequence analysis revealed that the BinA (41.9 kDa) and BinB (51.4kDa) proteins of KS2-13 and *L. sphaericus* 2362 differ by 3 (K89E, E104A, Y176D) and 6 (A69S, K70N, I110T, N248H, H314L and L317F) amino acids respectively. Similarly BinA and BinB proteins of KS2-15 and *L. sphaericus* 2362 strains vary by 1(E104A) and 3 (H109P, N248H and P274S) amino acids respectively. The varied amino acid sequences could be reason for the difference in activity. These two strains can act as good candidates for insecticidal formulation. Moreover, we reported for the first time the isolation of mosquitocidal *Lysinibacillus* strains from Kashmir valley.

Keywords: Bin proteins; *bin* genes; Kashmir valley; *Lysinibacillus*; SDS-PAGE.

Vector-borne diseases cause main public health tribulations and their control is largely accomplished with usage of synthetic insecticides against carrier insects (Baird 2000). Although the application of chemical insecticides proved lucrative to effectively control mosquitoes for many decades, the usage of chemical pesticides in

long term will result in malicious effect on human and environment. There are many health problems associated to pesticide usage which vary from abdominal pain, dizziness, headaches, nausea, vomiting, as well as skin and eye problems to cancer and developmental defects as well (Lorenz, 2009). Side effects on environment range from non-target organism killing (harmless insects, birds, amphibians and fishes) to increasing resistance to mosquito (Denholm *et al.* 2002). Effects on non-target organisms and concern about accumulation

* To whom all correspondence should be addressed.
Tel.: 0427-2345766; Fax: 0427-2345124;
E-mail: iarulselvibiotech@gmail.com

of chemicals in the environment have hastened the requirement to develop substitutes. Control of mosquitoes by means of entomopathogenic bacteria is a potential environmental friendly alternative to chemical insecticides (Park and Federici. 2009). *L. sphaericus* is an aerobic, endospore-forming gram-positive bacterium (Suryadi *et al.* 2016). This bacterium shows high toxicity against mosquito larvae and has been used extensively in some countries as biopesticide (Poopathi and Abidha, 2010). The larvicidal properties of this bacterium are mainly attributed to the presence of Binary proteins (BinA 41.9 kDa and BinB 51.4 kDa) expressed during sporulation stage (Broadwell *et al.* 1990; Berry, 2012). Cry48/Cry49 toxin expressed during sporulation in some strains play significant in the activity against *Culex* mosquitoes (Jones *et al.* 2007) and Mtx proteins (100-kDa toxin) formed during vegetative growth (Priest *et al.* 1997; Wirth *et al.* 2014). However Bin toxins are the key factors responsible for larvicidal activity. These Bin toxins after intake by susceptible larvae dissolve in the alkaline midgut and get activated by gut proteases. The 41.9 kDa BinA protein is cleaved to 39 kDa, and 51.4 kDa BinB is changed to 43 kDa (Baumann *et al.* 1991). BinB binds to the specific receptors present on larval midgut brush border membranes (Silva-Filha *et al.* 1999), while activated BinA brings the toxicity by working together with BinB (Oei *et al.* 1992; Lekakarn *et al.* 2015).

Valley Kashmir that is often referred as Terrestrial Paradise on Earth is situated at northern western tip of Himalayan biodiversity hot spot (Mittermeier *et al.*, 2005). It is located roughly between 32°15'2" and 37°05'2" North latitude and 72°35'2" to 83°20'2" East longitude, with complicated geomorphologic characteristics (i.e. snow clad mountains, vast meadows full of flowers, thick forests, small mountains, valley lakes and numerous serpentine rivers). These distinctive features associated with variety of animal forms ranging from higher groups such as vertebrates, including mammals, birds, reptiles, amphibians to lower groups like invertebrates including insects and even unicellular micro-organisms provide the opportunity to isolate novel mosquitocidal bacterial strains. In this study, we reported isolation and characterization of two highly mosquitocidal *B. sphaericus* strains from Himalayan valley Kashmir,

active against larvae of *C. quinquefasciatus* and *A. stephensi* epidemiological disease causing vectors.

MATERIALS AND METHODS

Reference strain

L. sphaericus 2362 was obtained from *Bacillus* Genetic Stock Center (Columbus, Ohio) which served as positive control.

Mosquito cultures

The cultures of *Culex quinquefasciatus*, *A. stephensi* and *A. aegypti* were procured from the Centre for Research in Medical Entomology (CRME), Madurai, Tamil Nadu and maintained at 28±2 °C and 75% to 85% relative humidity under a photoperiod of 14L:10D in our laboratory. Larvae were reared in chlorine free water and fed with dog yeast and biscuits at 2:3 ratio. For all bioassays, third-instar larvae of the size and same age were used.

Sample collection

A total of 158 samples from different divisions (Anantnag, Kulgam, Pulwama, Shopian, Budgam, Srinagar, Ganderbal, Bandipora, Baramulla, Kupwara) of the Himalayan valley Kashmir, India, were used for isolation of mosquitocidal bacteria. To our best knowledge, microbial insecticides had not been formerly applied in the sampled areas. All the soil samples (each ~ 5 g) were collected from 2 to 4 cm below the surface after scrapping off the surface material with a sterile spatula.

Isolation of Bacteria

Isolation of bacterial strains from the soil samples was performed according to the method described by Geetha *et al.* (2007) with slight modification. In the laboratory, One gram of soil sample was suspended in 10 mL of sterile distilled water (10⁻¹) in a boiling tube. One ml from this suspension was added to nine ml of water which gives 10⁻² dilution. Similarly, dilutions were made up to 10⁻⁵ and 0.1 ml from each dilution was spread on nutrient yeast salt mineral agar (NYSM) comprising of 5 g peptone, 5 g yeast extract, 3 g beef extract, 5 g NaCl, 5 g glucose, 103 mg CaCl₂, 10 mg MnCl₂ and 203 mg MgCl₂, (Hi-Media, India) per liter of distilled water. The bacterial suspensions were subjected to pasteurization before plating, expecting only gram-positive bacteria. The bacterial colonies which appeared on NYSM plates

after incubation at 30 °C for 48 h and were further purified using NYSM agar. Purified colonies were then sub-cultured on NYSM agar slants and stored at 4 °C. These bacterial isolates were tested for mosquito larvicidal activity.

Determination of larvicidal activity

A loopful of bacterial culture from each NYSM medium slant was inoculated to 3 ml of NYSM broth. This was incubated in a shaking incubator (Hasthas Scientific Instruments, India) maintained at 30 °C and 200 rpm till e⁹⁰% sporulation (472 h). After incubation, 1 ml (about 10⁹ spores) from the whole culture was utilized to screen for mosquito larvicidal activity. Bioassays with bacterial suspension (1 mL) were carried out in wax coated paper cups each containing 25 third instar larvae and 25 pupae of *C. quinquefasciatus*/*A. stephensi*/*A. aegypti* in 125 ml chlorine-free tap water at a temperature of 28±2 °C and 75% to 85% relative humidity under a photoperiod of 14L:10D. In control cups 1 ml of un-inoculated NYSM broth only was used and in another control 1ml of *L. sphaericus* 2362 was added. Mortality in the individual cups was noted down by counting the number of live larvae or pupae present subsequent to 24 h of culture introduction. Based on the mosquitocidal activity two strains were selected, code named as KS2-13 and KS2-15 and further characterized. Further bioassays with larvicidal active isolates (KS2-13 and KS2-15) and *L. sphaericus* 2362 were performed using seven different concentrations (10², 10³, 10⁴, 10⁵, 10⁶, 10⁷ and 10⁸) of bacterial spore crystal suspension obtained by serial dilution. All the bioassays were performed in ten replicates in wax coated paper cups each containing 25 third instar larvae of *C. quinquefasciatus*/*A. stephensi* in 125 ml chlorine-free tap water at 28±2 °C and 75% to 85% relative humidity under a photoperiod of 14L:10D. A control group, tested on NYSM broth without bacterial suspension, was included in each experiment. No source of food was added during bioassays. The number of dead larvae in each cup was counted after 24 h of culture introduction. Probit regression analysis was performed with SPSS 20.0 for windows software and LC50 and LC90 as well as their 95% fiducial limits were determined (Geetha *et al.* 2007; Tranchida *et al.* 2011).

Characterization of the potential bacterial strains

Two potential strains were studied for morphological, biochemical and physiological characteristics according to Bergey's Manual of Systematic Bacteriology (Sneath, 1986). Biochemical tests namely, fermentation of arabinose, glucose, mannitol, xylose and malonate, hydrolysis of starch, decarboxylation of lysine, ornithine, arginine dihydrolase, utilization of citrate, degradation of tyrosine, deamination of phenylalanine, nitrate reduction, decomposition of urea, indole and acetyl methyl carbinol production were performed. The capability to grow on different concentrations (2%, 5%, 7% and 10%) of NaCl was also studied.

Molecular characterization of the mosquitocidal bacterial strains

Genomic DNA was extracted from KS2-13 and KS2-15 using HiPurA™ Bacterial and Yeast Genomic DNA Purification Spin Kit (Hi-Media, India). To determine the sequence of the 16S rRNA gene, a DNA fragment of ~ 1.5-kb was amplified by PCR from the genomic DNA of the samples using universal eubacteria-specific primers: 27F (5'CTGAGAGTTTGATCMTGGCTCAG-3'CTG) and 1492R (5'GGYTACCTTGTTACGACTT-3'CTG), synthesized at Xcelris Labs Ltd, Gujarat, India. For polymerase chain reactions (PCR), 0.1 µg of total DNA from each isolate was mixed with 10 µl of 2X PCR Master Mix (GeNei™ Bengaluru, India) consisting of dNTPs, Taq polymerase and PCR buffer. Forward and reverse primers were used at a concentration of 1 µM. The final volume was made upto 20 µl with sterile double distilled water. PCR amplification was performed in a thermal cycler (cyber cycler-P series PCR peltier model p96+ USA) using the program: a 5 min denaturation step at 94 °C, 30 amplification cycles of 1 min at 94 °C, 1 min at 56 °C, and 1 min at 72 °C, with a final extension step of 10 min at 72 °C. The amplified PCR products were purified using GeneJET™ PCR purification kit (Fermentas life science Mumbai India) and sequenced by automated sequencer (ABI 3730xl Genetic) at Xcelris Labs Ltd, Gujarat, India. The forward and reverse sequences were edited using Bioedit program (Hall, 1999). The sequences obtained from strains KS2-13 and KS2-15 were compared to 16S ribosomal DNA (rDNA) gene

sequences available in the databases of the National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov/>) by BLASTN homology search, as described by Altschul *et al.* (1997). Phylogenetic analysis included the 16S rDNA gene sequences of the local isolates KS2-13 and KS2-15 and the reference strains of *L. sphaericus*, and *L. fusiformis* obtained from GenBank. *Alycyclobacillus cycloheptanicus* was used as an outgroup (Nakamura. 2000). Neighbor-Joining method (Saitou and Nei. 1987) was used to infer the evolutionary history. Cluster support was assessed through 1,000 bootstrap replicates (Felsenstein. 1985). The branch lengths of the drawn tree were in the similar units as those of the evolutionary distances used to deduce the phylogenetic tree. A Kimura 2-parameter method was used to compute the evolutionary distances and is in the units of the number of base substitutions per site (Kimura. 1980). The analysis entailed 17 nucleotide sequences. All positions having missing data and gaps were removed. There were a total of 1366 nucleotides positions in the final dataset. Phylogenetic and Evolutionary analyses were conducted in MEGA5 (Tamura *et al.* 2011). Sequences were submitted in GenBank and accession numbers were obtained.

SDS PAGE analysis of spore crystal mixture

Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) of spore crystal mixture was carried out by the method of Laemmli (1970) using 10% running and 4% stacking gels. The gels were stained with 0.4% Coomassie blue R250. The molecular mass of proteins was determined by using higher range protein molecular weight marker (myosin rabbit muscle 205kDa, phosphorylase b 97.4kDa, bovine serum albumin 66kDa, ovalbumin 43kDa and carbonic anhydrase 29kDa) obtained from GeNei™ Bangalore, India.

Screening for the presence of *L. sphaericus* toxin genes

Total cellular DNA isolated from indigenous *L. sphaericus* strain (KS2-13 and KS2-15) was used for identification of the dipterans-specific binary genes. Primers (Table1) designed by Hire *et al.* (2010) for the detection of binary toxins was used. The PCR amplification was performed in a thermal cycler (cyber cycler-P series PCR peltier model p96+ USA) in a 20µL reaction volume containing 100 ng DNA, 0.5 mM of primers,

2X PCR Master Mix consisting of dNTPs, Taq polymerase and PCR buffer (GeNei™ Bangalore, India). The amplified PCR products were purified using GeneJET™ PCR purification kit (Fermentas life science, Mumbai India).

Cloning and sequencing of binary genes

InsTAclone™ PCR Cloning Kit (Fermentas Life Science, Mumbai India) was used for cloning of purified PCR products. The PCR amplified *binA* and *binB* sequences were ligated in pTZ57R/T vector as per instructions in user manual. The recombinant vectors pTZ57R/T-*binA* and pTZ57R/T-*binB* were transformed into *Escherichia coli* DH5α. Positive clones were identified by blue white screening. The nucleotide sequences of two positive clones each from the pTZ57R/T-*binA* and pTZ57R/T-*binB* constructs were confirmed by complete sequencing of *binA* and *binB* using an automated DNA sequencer (ABI 3730xl Genetic) at Xcelris Labs Ltd, Gujarat, India. The forward and reverse sequences were edited using Bioedit program (Hall. 1999) and blast performed using BLASTN. Sequences were submitted in GenBank and accession numbers were obtained.

Comparative analysis of deduced amino acid

The amino acid sequences were deduced from the complete DNA nucleotide coding sequences of binary genes (KS2-13 and KS2-15) using online ExPASy translation tool. These were compared with the sequences of BinA and BinB proteins of reference strain *L. sphaericus* 2362 by Clustal Omega (www.ebi.ac.uk/Tools/msa/clustalo/) an online multiple sequence alignment program for amino acid variations.

RESULT

A total of 450 bacteria from 158 samples were selected randomly and screened for mosquitocidal activity. Preliminary screening with 1 ml of culture demonstrated that two bacterial strains isolated from soil samples have mosquito larvicidal properties with 100% mortality towards *C. quinquefasciatus* and *A. stephensi* after 24 hours of exposure. However, none of the isolate showed any activity against pupae of mosquito cultures. Subsequent bioassays of these two strains against larvae of *C. quinquefasciatus* and *A. stephensi* with different culture concentrations obtained by serial

dilutions showed promising activities at very low concentrations (Table 2) which were comparable to control *L. sphaericus* 2362 strain. These strains, code named as KS2-13 and KS2-15 were selected and studied further.

Characterisation of the potential mosquitocidal bacterial strains

Both the mosquitocidal bacterial isolates were found to be gram-positive, aerobic, motile, rod-shaped bacteria with terminal spherical sporangium. Table 3 summarizes results of the morphological and biochemical tests performed. On the basis of morphological and physiological characters, strains KS2-13 and KS2-15 were identified as *L. sphaericus*. PCR amplification of the 16 S rRNA genes of KS2-13 and KS2-15 yielded amplicon of ~1.5 kb size. BLASTN (NCBI) analyses of KS2-13 sequence indicated

a 99% similarity with other *L. sphaericus* strains (e.g., strain C5, accession no. KF523303.1; strain Ot4b.39, accession no. JQ744626.1 and strain VCRC B543, accession no. JN377786.1). Similarly, BLASTN result of KS2-15 also showed a 99% similarity with other *L. sphaericus* strains (e.g. strain Y73, accession no. JX067902.1; strain C5, accession no. KF523303.1 strain Ot4b.39, accession no. JQ744626.1). The 16S rRNA genes of KS2-13 and KS2-15 were submitted in GenBank under accession numbers KJ183073.1 and KJ183074.1 respectively. Next, we determined the degree of relatedness of our isolates to different *L. sphaericus* and *L. fusiformis* species through a phylogenetic analysis. The NJ tree shows a close relationship between the strains isolated in the present study and Group I of Nakamura (strains B-23269 and B-23287) (Fig. 1).




Fig. 1. Phylogenetic relationships are based on 16S rRNA gene sequence analysis of members of local isolates (KS2-13 and KS2-15) and the *Lysinibacillus* species. Evolutionary distances were calculated using the Kimura two parameters method, and the topology was inferred using the neighbor joining (NJ) method. Numbers above branches represent percentage bootstrap values based on 1,000 replicates. The 16S rRNA gene sequences of *Bacillus cycloheptanicus* was arbitrarily chosen as the out group. Accession numbers are between parentheses

SDS -PAGE analysis

Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) of spore mixture revealed two major polypeptide chains of approximately 51.4 kDa and 41.9 kDa which correspond to known consistent sizes of BinA and BinB proteins (Fig. 2).

Screening for the presence of *L. sphaericus* toxin genes

Primers for *binA* and *binB* gave rise to fragments of 1.1 kb and 1.3 kb amplicons, corresponding to the *binA* toxin (41.9 kDa) gene and *binB* toxin (51.4 kDa) gene (Fig. 3) respectively. The sequences of *binA* and *binB* genes have showed 99% and 100% identity with the same genes found in *L. sphaericus*. The sequences were submitted to the GenBank under accession no. KJ547612, KJ547613, KJ547614 and KJ547615.

Comparison of deduced amino acid sequences

The deduced amino acid sequences of *binA* and *binB* genes of KS2-13 and KS2-15 strains when compared with deduced amino acid sequences of *binA* and *binB* genes of *L. sphaericus* 2362 revealed that the BinA (41.9 kDa) and BinB (51.4kDa) proteins of KS2-13 and *L. sphaericus* 2362 differ by 3 (K89E, E104A, Y176D) and 6 (A69S, K70N, I110T, N248H, H314L and L317F) amino acids respectively. Similarly BinA and BinB

proteins of KS2-15 and *L. sphaericus* 2362 strains vary by 1 (E104A) and 3 (H109P, N248H and P274S) amino acids respectively. The varied amino acid sequences may be the reason for difference in activity.

DISCUSSION

In the course of diverse microbial pesticides, *Bacillus thuringiensis* and *L. sphaericus* are the most widely used alternative control agents for mosquitoes (Geetha *et al.* 2007; Hayes *et al.* 2011; Prabhu *et al.* 2013). *B. sphaericus* strains have been isolated worldwide from diverse habitats, including the soil (Radhika *et al.* 2011; Suryadi *et al.* 2016), aquatic habitats (Foda *et al.* 2013), excreta of arid birds (Poopathi *et al.* 2014) and dead larvae (Tranchida *et al.* 2011). In this study, for the first time, we have reported the isolation and characterization of the mosquitocidal strains from soil samples of Kashmir valley that is situated at northern western tip of Himalayan biodiversity hotspot (Mittermeier *et al.* 2005). A total of 450 bacterial isolates from 158 samples were screened for mosquitocidal activity. However, only two isolates code named as KS2-13 and KS2-15, being toxic were further studied. Level of




Fig. 2. SDS-PAGE analysis of spore crystal mixture of *L. sphaericus* strains: Lane M: High molecular weight marker; Lane 1-2: *L. sphaericus* strains, KS2-13 and KS2-15 respectively




Fig. 3. PCR amplification of *binA* and *binB* genes from *L. sphaericus* strains KS2-13 and KS2-15. Lane M, molecular weight marker; Lane C1 and C2: negative controls; Lane 1-2: *binA* genes (KS2-13 and KS2-15 strain); and Lane 3-4, *binB* genes (KS2-13 and KS2-15 strain)

toxicity was higher in case of *C. quinquefasciatus* than *Anopheles stephensi*. The isolate KS2-13 and KS2-15 did not produce mortality in *A. aegypti* larvae. Earlier it had been reported that fluorescent labeled *L. sphaericus* toxin binds proficiently with cultured *C. quinquefasciatus* cells and larval midgut cells in contrast to *A. aegypti* cultured cells and larval midgut cells where it binds slightly or is undetectable (Davidson *et al.* 1987; Davidson 1988). However, Tranchida *et al.* (2011) have reported isolation of *Lysinibacillus* strains from *Culex pipiens* larvae active against larvae of *Culex*, *Aedes*, *Culex*, *Ochlerotatus*, and *Anopheles* species. Identification of *B. sphaericus* like organisms is hard and arduous as they cannot be differentiated from each other by conventional phenotypic tests. The sequence of 16S rRNA genes has been widely used to identify an unknown bacterium to the genus or species level (Geetha *et al.* 2008). In present study, two potential mosquito larvicidal toxic strains were identified as *B. sphaericus* by their morphological, biochemical features and 16S rRNA gene sequence. Blast analysis of 16S rRNA gene sequences of KS2-13 and KS2-15 showed 99% identity with other already reported *B. sphaericus*. A typical feature used in the depiction of the *L. sphaericus* species is the lack of ability to

utilize pentoses and hexoses as the solitary carbon source. This is due to the absence of the *pgi* genes encoding enzymes for breaking and transporting these sugars (Hu *et al.* 2008). Our two isolates KS2-15 and KS2-13 exhibited promising mosquito larvicidal activity against *C. quinquefasciatus* (LC₅₀: 1.36×10^3 spores/mL; 1.41×10^3 spores/mL respectively) and *A. stephensi* (LC₅₀: 2.14×10^3 spores/mL; 2.11×10^3 spores/mL correspondingly) which was comparable to control *L. sphaericus* 2362 (*C. quinquefasciatus* LC₅₀: 1.34×10^3 spores/mL; *A. stephensi* 2.10×10^3 spores/mL) under our laboratory conditions. Tranchida *et al.* (2011) reported two *B. sphaericus* strains C107 and C207 that had showed highest mosquitocidal activity against *Culex pipiens* and *Ochlerotatus albifasciatus* with LC₅₀: 4×10^4 spores/mL and LC₅₀ of 3.4×10^6 spores/mL respectively. Recently, Suryadi *et al.* (2016) reported four mosquitocidal *L. sphaericus* strains MNT, SKT, TJJ2, and SLG. *L. sphaericus* strain MNT showed LC₅₀ of 3.70×10^5 cell/mL, 1.76×10^7 cell/mL and 4.45×10^7 cell/mL against *Culex*, *Anopheles* and *Aedes* respectively. Similarly strains SKT, TJJ2 and SLG showed LC₅₀ of 1.13×10^5 cell/mL, 2.85×10^5 cell/mL, 1.78×10^7 cell/mL; 1.03×10^5 cell/mL, 8.94×10^4 cell/mL, 1.72×10^7 cell/mL and 9.41×10^5 cell/mL, 2.39

Table 1. Primers used in this study

Gene	Primer sequence (5'E§ to 3'E§)	Size	Reference
<i>binA</i> gene	F= AGC TAA AAC ATATGA GAA ATT TGG ATT TTA TTG R= TTG TGG ATCCTTAGT TTT GAT CAT CTG TAATAATC	1.1 kb	Hire <i>et al.</i> , 2010
<i>binB</i> gene	F= GAT GAA GAA CATATG TGC GAT TCA AAA GAC R= AGT TGG ATC CTT ACT GGT TAA TTT TAG GTA TTA A	1.3 kb	

Table 2. Larvicidal activity of *Lysinibacillus sphaericus* strain KS2-13 and KS2-15 against third instar larvae of different mosquito species

Strain	Mosquito species	LC50 (95% CL)	LC90 (95% CL)	X ²
2362	<i>Culex quinquefasciatus</i>	1.34 (1.09-1.56)	2.48 (2.15- 3.08)	0.343
KS2-13	<i>Culex quinquefasciatus</i>	1.41 (1.14-1.65)	2.67 (2.29- 3.41)	0.190
KS2-15	<i>Culex quinquefasciatus</i>	1.36(1.13-1.58)	2.47 (2.15- 3.05)	0.313
2362	<i>Anopheles stephensi</i>	2.10 (1.78-2.69)	3.77 (3.04-5.70)	0.543
KS2-13	<i>Anopheles stephensi</i>	2.11 (1.74-2.91)	4.11 (3.18-7.06)	0.649
KS2-15	<i>Anopheles stephensi</i>	2.14 (1.81-2.78)	3.86 (3.09-5.94)	0.729

Lethal concentrations (expressed in 10^3 spores mL⁻¹) for 50% (LC₅₀) or 95% (LC₉₅) of larvae treated after 24 h

$\times 10^5$ cell/mL, 2.08×10^7 cell/mL against *Culex*, *Anopheles* and *Aedes* respectively. The toxicity of *L. sphaericus* to mosquito larvae mainly results from binary toxins (41.9 kDa and 51.4 kDa) encoded by the *binA* and *binB* genes expressed during sporulation stage (Broadwell *et al.* 1990; Tangsongcharoen *et al.* 2015). Another toxin called Mtx (100 kDa,) is present in *L. sphaericus* strains of both low and high toxicity. According to Thanabalu *et al.* (1991), the existence of the *mtx* genes does not per se confer toxicity to this bacterial strain against mosquito larvae. Therefore, the low toxicity in some strains could result from either low expression or short lived stability of the

binary toxins during sporulation. Recently Prabhu *et al.* (2013) reported the molecular characterization of forty two *L. sphaericus* strains isolated from Tamil Nadu, India. Their results revealed genetic heterogeneity between both toxic and non-toxic isolates and pointed out there is a good correlation between the existence of toxin genes and toxicity of the strains. In current study SDS-PAGE analysis of spore crystal revealed two major protein bands of 41.9 kDa and 51.4 kDa in size which correspond to the BinA and BinB proteins. The PCR for *binA* and *binB* genes and their sequence analysis confirmed the presence of two *bin* genes in each strain. Deduced amino acid sequences of *binA* and

Table 3. Morphological and biochemical characteristics of mosquitocidal bacterial strains

Property	KS2-13	KS2-15
Colony morphology	Cream colored, moist irregular	Cream colored , irregular
Spores	Cylindrical	cylindrical
Utilization of		
Glucose	-	-
Starch	-	-
Lactose	-	-
arabinose	-	-
Xylose	-	-
Mannitol	-	-
2% NaCl	+	+
4% NaCl	+	+
5% NaCl	W	W
7% NaCl	-	-
Erythromycin 1b	+	+
Erythromycin 2b	-	-
Tetracycline 2b	+	+
Tetracycline 5b	-	-
Chloramphenicol 8b	+	+
Citrate	+	+
Phenyl alanine	+	+
Arginine	-	-
Lysine	-	-
Ornithine	-	-
Tyrosine	-	-
H ₂ S production	-	-
Decomposition of urea	W	W
Catalase	+	+
Oxidase	+	+
Indole	-	-
Methyl red	-	-
Voges-Proskauer	+	+
Identified as	<i>Lysinibacillus sphaericus</i>	<i>Lysinibacillus sphaericus</i>

+ indicates positive; - indicates negative and W indicates weak

binB genes from KS2-13 and KS2-15 strains when compared with amino acid sequences of *binA* and *binB* genes from *L. sphaericus* 2362 revealed that the BinA and BinB proteins of KS2-13 and *L. sphaericus* 2362 are different by 3 (K89E, E104A, Y176D) and 6 (A69S, K70N, I110T, N248H, H314L and L317F) amino acids respectively. Similarly BinA and BinB proteins of KS2-15 and *L. sphaericus* 2362 strains contrast by 1 (E104A) and 3 (H109P, N248H and P274S) amino acids respectively. The varied amino acid sequences may be reason for the difference in activity.

The high larval toxicity existing in some *L. sphaericus* strains, such as 2362 (Weiser, 1984) and IAB59 (de Barjac *et al.* 1988), has resulted into their commercial use as biopesticides against populations of mosquitos. Recent reports of resistance development devoted the attention of researchers worldwide towards the discovery of new isolates from natural resources as an alternative to the existing biopesticides globally. As variability was found between the new isolated *L. sphaericus* strains as well as with *L. sphaericus* strains 2362, we concluded that the strains (KS2-13 and KS2-15) isolated and identified in this study embody good candidates for exploiting in the control of mosquito within the situation to which both the host and the parasite are evenly well adapted.

ACKNOWLEDGEMENT

We would like to thank Dr. Daniel R. Zeigler (Ph.D), director of Bacillus Genetic Stock Center (Columbus, Ohio) for providing *L. sphaericus* 2362. We also extend our thanks to Director of 'Centre for Research in Medical Entomology (CRME)' Madurai, Tamil Nadu for providing mosquito cultures: *Culex quinquefasciatus*, *A. stephensi* and *A. aegypti*.

REFERENCES

- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J., Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*, 1997; **25**(17):3389-63402.
- Baird, J.K., Resurgent malaria at the millennium: control strategies in crisis. *Drugs*, 2000; **59**(4):719-743.
- Baumann, P., Clark, M.A., Baumann, L., Broadwell, A.H., *Bacillus sphaericus* as a mosquito pathogen: properties of the organism and its toxins. *Microbiol Rev*, 1991; **55**(3): 425-6436.
- Berry, C. The bacterium, *Lysinibacillus sphaericus*, as an insect pathogen. *Journal of Invertebrate Pathology*, 2012; **109**: 1-10
- Broadwell, A.H., Baumann, L., Baumann, P., The 42- and 51- kilodalton mosquitocidal proteins of *Bacillus sphaericus* 2362: construction of recombinants with enhanced expression and in vivo studies of processing and toxicity. *J Bacteriol*, 1990; **172**(5):2217-2223.
- Davidson, E.W. Binding of the *Bacillus sphaericus* (Eubacteriales: Bacillaceae) toxin to midgut cells of mosquito (Diptera: Culicidae) larvae: relationship to host range. *J. Med Entomol*, 1988; **25**(3):151-6157.
- Davidson, E.W., Shellabarger, C., Meyer, M., Bieber, A.L., Binding of the *Bacillus sphaericus* mosquito larvicidal toxin to cultured insect cells. *Canad. J. Microbiol*, 1987; **33**(11): 982-6989.
- de Barjac, H., Thiery, I., Cosmao-Dumanoir, V., Frachon, E., Laurent, P., Charles, J.F., Hamon, S., Ofori, J., Another *Bacillus sphaericus* serotype harbouring strains very toxic to mosquito larvae: serotype H6. *Ann Inst Pasteur Microbiol*, 1988; **139**(3):363-6377.
- Denholm, I., Devine, G.J., Williamson, M.S., Evolutionary genetics. Insecticide resistance on the move. *Science*, 2002; **297**(5590): 2222-3.
- Felsenstein, J., Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 1985; **39**(4):783-6791.
- Foda M.S., Amin M. A, El Tayeb O. M., Gawdat N. A. and El Bendary M. A. Isolation and characterization of highly potent mosquitocidal bacillus from Egyptian environment. *J.Biological Sci*, 2013; **13**(6): 483-490
- Geetha, I., Manonmani, A.M. Mosquito pupicidal toxin production by *Bacillus subtilis* subsp. *subtilis*. *Biol Control*, 2008; **44**(2):242-6247.
- Geetha, I., Prabakaran, G., Paily, K.P., Manonmani, A.M., Balaraman, K., Characterisation of three mosquitocidal *Bacillus* strains isolated from mangrove forest. *Biol Control*, 2007; **42**(1): 34-640.
- Hall, T.A., BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser*, 1999; **41**:95-698.
- Hayes, S.R., Hudon, M, Park, H.W., Isolation of novel *Bacillus* species showing high mosquitocidal activity against several mosquito species. *J. Invertebr Pathol*, 2011; **107**(1):139-57660.

16. Hire, R.S., Hadapad, A.B., Vijayalakshmi, N., Dongre, T.K., Characterization of highly toxic indigenous strains of mosquitocidal organism *Bacillus sphaericus*. *FEMS Microbiol Lett*, 2010; **305**(2):155-6-161.
17. Hu, X., Fan, W., Han, B., Liu, H., Zheng, D., Li, Q., Dong, W., Yan, J., Gao, M., Berry, C., Yuan, Z., Complete genome sequence of the mosquitocidal bacterium *Bacillus sphaericus* C3-41 and comparison with those of closely related *Bacillus* species. *J Bacteriol.*, 2008; **190**(8):2892-6-2902.
18. Jones GW, Nielsen-Leroux C, Yang Y, Yuan Z, Dumas VF, Monnerat RG, Berry C. A new Cry toxin with a unique two-component dependency from *Bacillus sphaericus*. *FASEB J.*, 2007; **21**(14):4112-20.
19. Kimura, M., A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol*, 1980; **16**(2):111-6-120.
20. Laemmli, U.K., Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* (London), 1970; **227**(5259):680-6-685.
21. Lekakarn H., Promdonkoy B., Boonserm P. Interaction of *Lysinibacillus sphaericus* binary toxin with mosquito larval gut cells: Binding and internalization. *Journal of Invertebrate Pathology*, 2015; **132**: 125-131
22. Lorenz, E.S. Potential health effects of pesticides. Pennsylvania: Penn State's College of Agricultural Sciences 2009. [Online] Available from: http://extension.psu.edu/pests/pesticide-education/applicators/fact-sheets/pesticide-safety/potential-health-effects-of-pesticides/extension_publication_file [Accessed on 3rd March, 2016]
23. Mittermeier, R.A., Gil, P.R., Hoffmann, M., Pilrim, J., Brooks, T., Mittermeier, C.G., Lamoreux, J., Fornseca, G.A.B. Hotspots revisited: Earth's biologically richest and most endangered terrestrial eco-regions. Boston: University of Chicago, 2005; 392p.
24. Nakamura, L.K. Phylogeny of *Bacillus sphaericus*-like organisms. *Int J Syst Evol Microbiol.*, 2000; **50** Pt5: 1715-6-1722.
25. Oei, C., Hindley, J., Berry, C., Binding of purified *Bacillus sphaericus* binary toxin and its deletion derivatives to *Culex quinquefasciatus* gut: elucidation of functional binding domains. *J Gen Microbiol*, 1992; **138**(7): 1515-6-1526.
26. Park, H.W., Federici, B.A., Genetic engineering of bacteria to improve efficacy using the insecticidal proteins of *Bacillus* species. In: Stock, S.P. (Ed.), *Insect Pathogens: Molecular Approaches and Techniques*. *CABI International*, 2009; pp. 275-6-305.
27. Poopathi, S., Abidha, S. Mosquitocidal bacterial toxins (*Bacillus sphaericus* and *Bacillus thuringiensis* serovar israelensis): mode of action, cytopathological effects and mechanism of resistance. *Journal of Physiology and Pathophysiology*, 2010; **1**(3): 22-38.
28. Poopathi, S., Thirugnanasambantham, K., Mani, C., Ragul, K., Sundarapandian, S.M. Isolation of mosquitocidal bacteria (*Bacillus thuringiensis*, *B. sphaericus* and *B. cereus*) from excreta of arid birds. *Indian J Exp Biol.*, 2014; **52**(7):739-47.
29. Prabhua, D.I.G., Sankara, S.G., Vasan, P.T., Piriya, P.S., Selvana, B.K., Vennisona, S.J., Molecular characterization of mosquitocidal *Bacillus sphaericus* isolated from Tamil Nadu, India. *Acta Trop*, 2013; **127**(3):158-64.
30. Priest, F.G., Ebdrup, L., Zahner, V., Carter, P., Distribution and characterization of mosquitocidal toxin genes in some strains of *Bacillus sphaericus*. *Appl Environ Microbiol*, 1997; **63**(4):1195-6-1198.
31. Radhika, D., Ramathilaga, A., Prabu, C.S., Murugesan, A.G. Evaluation of larvicidal activity of soil microbial isolates (*Bacillus* and *Acinetobacter* Sp.) against *Aedes aegypti* (Diptera: Culicidae) - the vector of Chikungunya and Dengue. *Proceedings of the International Academy of Ecology and Environmental Sciences*, 2011; **1**(3-4):169-178
32. Saitou, N., Nei, M., The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol*, 1987; **4**(4): 406-6-425.
33. Silva-Filha, M.H., Nielsen-LeRoux, C., Charles J.F., Identification of the receptor for *Bacillus sphaericus* crystal toxin in the brush border membrane of the mosquito *Culex pipiens* (Diptera: Culicidae). *Insect Biochem Mol Biol*, 1999; **29**(8): 711-6-721.
34. Sneath, P.H.A., Endospore forming gram positive rods, cocci. In: Sneath, P.H.A., Mair N., Sharpe M, Holt J (Eds.), *Bergey's manual of systematic bacteriology*, vol. 2. Williams & Wilkins, Baltimore, MD, 1986; pp. 1104-6-1207.
35. Suryadi, B.F., Yanuwadi, B., Ardyati, T., Suharjo, S. Evaluation of entomopathogenic *Bacillus sphaericus* isolated from Lombok beach area against mosquito larvae. *Asian Pac J Trop Biomed*; 2016; **6**(2): 148-154
36. Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol Biol Evol*,

- 2011; **28**(10): 2731-6-2739.
37. Tangsongcharoen, C., Chomanee, N., Promdonkoy, B., Boonserm, P. *Lysinibacillus sphaericus* binary toxin induces apoptosis in susceptible *Culex quinquefasciatus* larvae. *J Invertebr Pathol.*, 2015; **128**:57-63
38. Thanabalu, T., Hindley, J., Jackson-Yap, J., Berry, C., Cloning, sequencing, and expression of a gene encoding a 100-kilodalton mosquitocidal toxin from *Bacillus sphaericus* SSII-1. *J Bacteriol.*, 1991; **173**(9):2776-2785.
39. Tranchida, M.C., Riccillo, P.M., Micieli, M.V., García, J.J., Rodríguez, M.S., Isolation, characterization and evaluation of mosquitocidal activity of *Lysinibacillus* strains obtained from *Culex pipiens* larvae. *Ann Microbiol.*, 2011; **61**(1): 575-6-584.
40. Weiser, J., A mosquito-virulent *Bacillus sphaericus* in adult *Simulium damnosum* from Northern Nigeria. *Zentralbl Microbiol* 1984 ; **139**(1):57-6-60.
41. Wirth, M.C., Berry, C., Walton, W. E., Federici, B. A. Mtx toxins from *Lysinibacillus sphaericus* enhance mosquitocidal cry-toxin activity and suppress cry-resistance in *Culex quinquefasciatus*. *Journal of Invertebrate Pathology*, 2014; **115**: 62-67